

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 September 2008 (18.09.2008)

PCT

(10) International Publication Number
WO 2008/110196 A1

(51) International Patent Classification:

C07D 403/06 (2006.01) A61P 3/10 (2006.01)
A61K 31/439 (2006.01) A61P 3/04 (2006.01)

(21) International Application Number:

PCT/EP2007/052219

(22) International Filing Date: 9 March 2007 (09.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **NOVO NORDISK A/S** [DK/DK]; DK-2880 Bagsvaerd (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KILBURN, John, Paul** [GB/DK]; DK-4690 Haslev (DK). **ANDERSEN, Henrik, Sune** [DK/DK]; DK-2800 Lyngby (DK). **KAMPEN, Gita, Camilla, Tejlgaard** [DK/DK]; DK-2850 Nærum (DK).

(74) Common Representative: **NOVO NORDISK A/S**; Novo Allé, DK-2880 Bagsvaerd (DK).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(54) Title: **INDOLE- AND BENZIMIDAZOLE AMIDES AS HYDROXYSTEROID DEHYDROGENASE INHIBITORS**



(I)

(57) Abstract: The use of substituted amides of structure (I) for modulating the activity of 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) and the use of these compounds as pharmaceutical compositions, are described. Also a novel class of substituted amides, their use in therapy, pharmaceutical compositions comprising the compounds, as well as their use in the manufacture of medicaments are described. The present compounds are modulators and more specifically inhibitors of the activity of 11 β HSD1 and may be useful in the treatment, prevention and/or prophylaxis of a range of medical disorders where a decreased intracellular concentration of active glucocorticoid is desirable.



WO 2008/110196 A1

INDOLE- AND BENZIMIDAZOLE AMIDES AS HYDROXYSTEROID DEHYDROGENASE INHIBITORS

FIELD OF INVENTION

[0001] The present invention relates to use of substituted amides and pharmaceutical compositions comprising the same for treating disorders where it is desirable to modulate the activity of 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1). The present invention also relates to novel substituted amides, to their use in therapy, to pharmaceutical compositions comprising the same, to the use of said compounds in the manufacture of medicaments, and to therapeutic methods comprising the administration of the compounds. The present compounds modulate the activity of 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) and are accordingly useful in the treatment of diseases in which such a modulation is beneficial, such as the metabolic syndrome.

BACKGROUND OF THE INVENTION

[0002] The metabolic syndrome is a major global health problem. In the US, the prevalence in the adult population is currently estimated to be approximately 25%, and it continues to increase both in the US and worldwide. The metabolic syndrome is characterised by a combination of insulin resistance, dyslipidemia, obesity and hypertension leading to increased morbidity and mortality of cardiovascular diseases. People with the metabolic syndrome are at increased risk of developing frank type 2 diabetes, the prevalence of which is equally escalating.

[0003] In type 2 diabetes, obesity and dyslipidemia are also highly prevalent and around 70% of people with type 2 diabetes additionally have hypertension once again leading to increased mortality of cardiovascular diseases.

[0004] In the clinical setting, it has long been known that glucocorticoids are able to induce all of the cardinal features of the metabolic syndrome and type 2 diabetes.

[0005] 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) catalyses the local generation of active glucocorticoid in several tissues and organs including predominantly the liver and adipose tissue, but also e.g. skeletal muscle, bone, pancreas, endothelium, ocular tissue and certain parts of the central nervous system. Thus, 11 β HSD1 serves as a local regulator of glucocorticoid actions in the tissues and organs where it is expressed (Tannin et al., *J. Biol. Chem.*, 266, 16653 (1991); Bujalska et al., *Endocrinology*, 140, 3188 (1999); Whorwood et al., *J Clin Endocrinol Metab.*, 86, 2296 (2001); Cooper et al., *Bone*, 27, 375 (2000); Davani et al., *J. Biol. Chem.*, 275, 34841 (2000); Brem et al., *Hypertension*, 31, 459 (1998); Rauz et al.,

Invest. Ophthalmol. Vis. Sci., 42, 2037 (2001); Moisan et al., *Endocrinology*, 127, 1450 (1990)).

[0006] The role of 11 β HSD1 in the metabolic syndrome and type 2 diabetes is supported by several lines of evidence. In humans, treatment with the non-specific 11 β HSD1 inhibitor carbenoxolone improves insulin sensitivity in lean healthy volunteers and people with type 2 diabetes. Likewise, 11 β HSD1 knock-out mice are resistant to insulin resistance induced by obesity and stress. Additionally, the knock-out mice present with an anti-atherogenic lipid profile of decreased VLDL triglycerides and increased HDL-cholesterol. Conversely, mice that overexpress 11 β HSD1 in adipocytes develop insulin resistance, hyperlipidemia and visceral obesity, a phenotype that resembles the human metabolic syndrome (Andrews et al., *J. Clin. Endocrinol. Metab.*, 88, 285 (2003); Walker et al., *J. Clin. Endocrinol. Metab.*, 80, 3155 (1995); Morton et al., *J. Biol. Chem.*, 276, 41293 (2001); Kotelevtsev et al., *Proc. Natl. Acad. Sci. USA*, 94, 14924 (1997); Masuzaki et al., *Science*, 294, 2166 (2001)).

[0007] The more mechanistic aspects of 11 β HSD1 modulation and thereby modulation of intracellular levels of active glucocorticoid have been investigated in several rodent models and different cellular systems. 11 β HSD1 promotes the features of the metabolic syndrome by increasing hepatic expression of the rate-limiting enzymes in gluconeogenesis, namely phosphoenolpyruvate carboxykinase and glucose-6-phosphatase, promoting the differentiation of preadipocytes into adipocytes thus facilitating obesity, directly and indirectly stimulating hepatic VLDL secretion, decreasing hepatic LDL uptake and increasing vessel contractility (Kotelevtsev et al., *Proc. Natl. Acad. Sci. USA*, 94, 14924(1997); Morton et al., *J. Biol. Chem.* 276, 41293 (2001); Bujalska et al., *Endocrinology*, 140, 3188 (1999); Souness et al., *Steroids*, 67, 195 (2002), Brindley & Salter, *Prog. Lipid Res.*, 30, 349 (1991)).

[0008] WO 01/90090, WO 01/90091, WO 01/90092, WO 01/90093 and WO 01/90094 disclose various thiazol-sulfonamides as inhibitors of the human 11 β -hydroxysteroid dehydrogenase type 1 enzyme, and further states that said compounds may be useful in treating diabetes, obesity, glaucoma, osteoporosis, cognitive disorders, immune disorders and depression.

[0009] We have now found substituted amides that modulate the activity of 11 β HSD1 leading to altered intracellular concentrations of active glucocorticoid. More specifically, the present compounds inhibit the activity of 11 β HSD1 leading to decreased intracellular concentrations of active glucocorticoid. Thus, the present compounds can be used to treat disorders where a decreased level of active intracellular glucocorticoid is desirable, such as e.g. the metabolic syndrome, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glu-

cose (IFG), dyslipidemia, obesity, hypertension, diabetic late complications, cardiovascular diseases, arteriosclerosis, atherosclerosis, myopathy, muscle wasting, osteoporosis, neurodegenerative and psychiatric disorders, and adverse effects of treatment or therapy with glucocorticoid receptor agonists.

- 5 **[0010]** One object of the present invention is to provide compounds, pharmaceutical compositions and use of compounds that modulate the activity of 11 β HSD1.

DEFINITIONS

- 10 **[0011]** In the following structural formulas and throughout the present specification, the following terms have the indicated meaning. The examples provided in the definitions present in this application are non-inclusive unless otherwise stated. They include but are not limited to the recited examples.

[0012] The term "halo" includes fluorine, chlorine, bromine, and iodine.

- 15 **[0013]** The term "trihalomethyl" includes trifluoromethyl, trichloromethyl, tribromomethyl, and triiodomethyl.

[0014] The term "trihalomethoxy" includes trifluoromethoxy, trichloromethoxy, tribromomethoxy, and triiodomethoxy.

- 20 **[0015]** The term "alkyl" includes C₁-C₈ straight chain saturated and methylene aliphatic hydrocarbon groups and C₃-C₈ branched saturated hydrocarbon groups having the specified number of carbon atoms. For example, this definition includes methyl (Me), ethyl (Et), propyl (Pr), butyl (Bu), pentyl, hexyl, isopropyl (i-Pr), isobutyl (i-Bu), *tert*-butyl (*t*-Bu), *sec*-butyl (*s*-Bu), isopentyl, and neopentyl.

- 25 **[0016]** The term "alkenyl" includes C₂-C₆ straight chain unsaturated aliphatic hydrocarbon groups and branched C₃-C₆ unsaturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, this definition includes ethenyl, propenyl, butenyl, pentenyl, hexenyl, methylpropenyl, and methylbutenyl.

- 30 **[0017]** The term "alkynyl" includes C₂-C₆ straight chain unsaturated aliphatic hydrocarbon groups and C₄-C₆ branched unsaturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, this definition includes ethynyl, propynyl, butynyl, pentynyl, hexynyl, and methylbutynyl.

- 35 **[0018]** The term "saturated or partially saturated monocyclic, bicyclic, or tricyclic ring system" represents but is not limited to aziridinyl, azepanyl, azocanyl, pyrrolinyl, pyrrolidinyl, 2-imidazoliny, imidazolidinyl, 2-pyrazoliny, morpholiny, piperidinyl, thiomorpholiny, piperaziny, phthalimide, 1,2,3,4-tetrahydro-quinoliny, 1,2,3,4-tetrahydro-isoquinoliny, 1,2,3,4-tetrahydro-quinoxaliny, indoliny, 1, 6-aza-bicyclo[3.2.1]octane, 2-aza-bicyclo[4.1.1]octane,

2-aza-bicyclo[3.2.1]octanyl, 7-aza-bicyclo[4.1.1]octanyl, 9-aza-bicyclo[3.3.2]decanyl, 4-aza-tricyclo[4.3.1.1^{3,8}]undecanyl, 9-aza-tricyclo[3.3.2.0^{3,7}]decanyl.

[0019] The term "saturated or partially saturated ring" represents cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclononenyl, cyclodecenyl, tetrahydrofuranyl, and tetrahydropyranyl.

[0020] The term "saturated or partially saturated aromatic ring" represents cyclopentyl, cyclohexyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclononenyl, cyclodecenyl, tetrahydrofuranyl, tetrahydropyranyl, phenyl, pyridyl, and pyrimidinyl.

[0021] The term "cycloalkyl" represents a saturated, mono-, bi-, tri- or spirocarbocyclic group having the specified number of carbon atoms (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, bicyclo[3.2.1]octyl, spiro[4.5]decyl, norpinyl, norbornyl, norcaryl, and adamantyl).

[0022] The term "cycloalkylalkyl" represents a cycloalkyl group as defined above attached through an alkyl group having the indicated number of carbon atoms or substituted alkyl group as defined above (e.g., cyclopropylmethyl, cyclobutylethyl, and adamantylmethyl).

[0023] The term "cycloalkenyl" represents a partially saturated, mono-, bi-, tri- or spirocarbocyclic group having the specified number of carbon atoms (e.g., cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclononenyl, and cyclodecenyl).

[0024] The term "cycloalkylcarbonyl" represents a cycloalkyl group as defined above having the indicated number of carbon atoms attached through a carbonyl group (e.g., cyclopropylcarbonyl and cyclohexylcarbonyl).

[0025] The term "cycloalkylalkylcarbonyl" represents a cycloalkyl group as defined above attached through an alkyl group having the indicated number of carbon atoms or substituted alkyl group as defined above (e.g., cyclohexylmethylcarbonyl and cycloheptylethylcarbonyl).

[0026] The term "hetcycloalkyl" represents a saturated mono-, bi-, tri-, or spirocarbocyclic group having the specified number of atoms with 1-4 of the specified number being heteroatoms or groups selected from nitrogen, oxygen, sulphur, SO, and SO₂ (e.g., tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidine, and pyridazine).

[0027] The term "hetcycloalkylalkyl" represents a hetcycloalkyl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., tetrahydrofuranylmethyl, tetrahydropyranylethyl, and tetrahydrothiopyranylmethyl).

[0028] The term "hetcycloalkylcarbonyl" represents a hetcycloalkyl group as defined above having the indicated number of carbon atoms attached through a carbonyl group (e.g., 1-piperidin-4-yl-carbonyl and 1-(1,2,3,4-tetrahydro-isoquinolin-6-yl)carbonyl).

[0029] The term "alkyloxy" represents an alkyl group having the indicated number of carbon atoms attached through an oxygen bridge (e.g., methoxy, ethoxy, propyloxy, allyloxy, and cyclohexyloxy).

[0030] The term "alkyloxyalkyl" represents an alkyloxy group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., methyloxymethyl).

[0031] The term "aryl" includes a carbocyclic aromatic ring that is monocyclic, bicyclic, or polycyclic, such as phenyl, biphenyl, naphthyl, anthracenyl, phenanthrenyl, fluorenyl, indenyl, pentalenyl, azulenyl, and biphenylenyl. Aryl also includes the partially hydrogenated derivatives of the carbocyclic aromatic enumerated above. Examples of partially hydrogenated derivatives include 1,2,3,4-tetrahydronaphthyl and 1,4-dihydronaphthyl.

[0032] The term "aryl1" includes phenyl, biphenyl, naphthyl, anthracenyl, phenanthrenyl, and fluorenyl.

[0033] The term "aryl2" includes phenyl, biphenyl, and naphthyl.

[0034] The term "hetaryl" includes pyrrolyl (2-pyrrolyl), pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), isoxazolyl (3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), thiophenyl (2-thiophenyl, 3-thiophenyl, 4-thiophenyl, 5-thiophenyl), furanyl (2-furanyl, 3-furanyl, 4-furanyl, 5-furanyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl), 5-tetrazolyl, pyrimidinyl (2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl), isoquinolyl (1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl), benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5-benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl)), 1,4-benzodioxin (2-(1,4-benzodioxin), 3-(1,4-benzodioxin), 5-(1,4-benzodioxin), 6-(1,4-benzodioxin)), benzo[b]thiophenyl (2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl, 7-benzo[b]thiophenyl), 2,3-dihydro-benzo[b]thiophenyl (2-(2,3-dihydro-benzo[b]thiophenyl), 3-(2,3-dihydro-benzo[b]thiophenyl), 4-(2,3-dihydro-benzo[b]thiophenyl), 5-(2,3-dihydro-benzo[b]thiophenyl), 6-(2,3-dihydro-benzo[b]thiophenyl), 7-(2,3-dihydro-benzo[b]thiophenyl)), 4,5,6,7-tetrahydro-benzo[b]thiophenyl (2-(4,5,6,7-tetrahydro-benzo[b]thiophenyl), 3-(4,5,6,7-tetrahydro-benzo[b]thiophenyl), 4-(4,5,6,7-tetrahydro-benzo[b]thiophenyl), 5-(4,5,6,7-tetrahydro-benzo[b]thiophenyl), 6-(4,5,6,7-tetrahydro-

benzo[b]thiophenyl), 7-(4,5,6,7-tetrahydro-benzo[b]thiophenyl)), thieno[2,3-b]thiophenyl, 4,5,6,7-tetrahydro-thieno[2,3-c]pyridyl (4-(4,5,6,7-tetrahydro-thieno[2,3-c]pyridyl), 5-4,5,6,7-tetrahydro-thieno[2,3-c]pyridyl), 6-(4,5,6,7-tetrahydro-thieno[2,3-c]pyridyl), 7-(4,5,6,7-tetrahydro-thieno[2,3-c]pyridyl)), indolyl (1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), isoindolyl (1-isoindolyl, 2-isoindolyl, 3-isoindolyl, 4-isoindolyl, 5-isoindolyl, 6-isoindolyl, 7-isoindolyl), 1,3-dihydro-isoindolyl (1-(1,3-dihydro-isoindolyl), 2-(1,3-dihydro-isoindolyl), 3-(1,3-dihydro-isoindolyl), 4-(1,3-dihydro-isoindolyl), 5-(1,3-dihydro-isoindolyl), 6-(1,3-dihydro-isoindolyl), 7-(1,3-dihydro-isoindolyl)), indazole (1-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6-benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (1-benz-oxazolyl, 2-benzoxazolyl), benzothiazolyl (1-benzothiazolyl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), benzo-[1,2,5]oxadiazolyl, (4-benzo[1,2,5]oxadiazole, 5-benzo[1,2,5]oxadiazole), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), piperidinyl (2-piperidinyl, 3-piperidinyl, 4-piperidinyl), and pyrrolidinyl (1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl).

[0035] The term "arylalkyl" represents an aryl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., benzyl, phenylethyl, 3-phenylpropyl, 1-naphtylmethyl, and 2-(1-naphtyl)ethyl).

[0036] The term "hetarylalkyl" or "hetaralkyl" represents a hetaryl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., (2-furyl)methyl, (3-furyl)methyl, (2-thienyl)methyl, (3-thienyl)methyl, (2-pyridyl)methyl, and 1-methyl-1-(2-pyrimidyl)ethyl).

[0037] The term "aryloxyhetaryl" represents an aryloxy group as defined above attached through a hetaryl group (e.g., 2-phenoxy-pyridyl).

[0038] The term "aryloxy" represents an aryl group as defined above attached through an oxygen bridge (e.g., phenoxy and naphthyloxy).

[0039] The term "hetaryloxy" represents a hetaryl group as defined above attached through an oxygen bridge (e.g., 2-pyridyloxy).

[0040] The term "arylalkyloxy" represents an arylalkyl group as defined above attached through an oxygen bridge (e.g., phenethyloxy and naphthylmethyloxy).

[0041] The term "hetarylalkyloxy" represents a hetarylalkyl group as defined above attached through an oxygen bridge (e.g., 2-pyridylmethyloxy).

[0042] The term "alkyloxycarbonyl" represents an alkyloxy group as defined above attached through a carbonyl group (e.g., methylformiat and ethylformiat).

[0043] The term "aryloxycarbonyl" represents an aryloxy group as defined above attached through a carbonyl group (e.g., phenylformiat and 2-thiazolylformiat).

[0044] The term "arylalkyloxycarbonyl" represents an "arylalkyloxy" group as defined above attached through a carbonyl group (e.g., benzylformiat and phenyletylformiat).

5 [0045] The term "alkylthio" represents an alkyl group having the indicated number of carbon atoms attached through a sulphur bridge (e.g., methylthio and ethylthio).

[0046] The term "arylthio" represents an aryl group as defined above attached through a sulphur bridge (e.g., benzenethiol and naphthylthiol).

10 [0047] The term "hetarylthio" represents a hetaryl group as defined above attached through a sulphur bridge (e.g., pyridine-2-thiol and thiazole-2-thiol).

[0048] The term "arylthioalkyl" represents an arylthio group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., methylsulfanyl benzene, and ethylsulfanyl naphthalene).

15 [0049] The term "hetarylthioalkyl" represents a hetarylthio group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., 2-methylsulfanyl-pyridine and 1-ethylsulfanyl-isoquinoline).

[0050] The term "hetaryloxyaryl" represents a hetaryloxy group as defined above attached through an aryl group as defined above (e.g., 1-phenoxy-isoquinolyl and 2-phenoxy-pyridyl).

20 [0051] The term "hetaryloxyhetaryl" represents a hetaryloxy group as defined above attached through a hetaryl group as defined above (e.g., 1-(2-pyridyloxy-isoquinoline) and 2-(imidazol-2-yloxy-pyridine)).

[0052] The term "aryloxyalkyl" represents an aryloxy group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., phoxymethyl and naphthyloxyethyl).

25 [0053] The term "aryloxyaryl" represents an aryloxy group as defined above attached through an aryl group as defined above (e.g., 1-phenoxy-naphthalene and phenyloxyphenyl).

[0054] The term "arylalkyloxyalkyl" represents an arylalkyloxy group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., ethoxymethyl-benzene and 2-methoxymethyl-naphthalene).

30 [0055] The term "hetaryloxyalkyl" represents a hetaryloxy group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., 2-pyridyloxymethyl and 2-quinolyloxyethyl).

35 [0056] The term "hetarylalkyloxyalkyl" represents a hetarylalkyloxy group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., 4-methoxymethyl-pyrimidine and 2-methoxymethyl-quinoline).

[0057] The term "alkylcarbonyl" represents an alkyl group as defined above having the indicated number of carbon atoms attached through a carbonyl group (e.g., octylcarbonyl, pentylcarbonyl, and 3-hexenylcarbonyl).

5 [0058] The term "arylcarbonyl" represents an aryl group as defined above attached through a carbonyl group (e.g., benzoyl).

[0059] The term "hetarylcarbonyl" represents a hetaryl group as defined above attached through a carbonyl group (e.g., 2-thiophenylcarbonyl, 3-methoxy-anthrylcarbonyl, and oxazolylcarbonyl).

10 [0060] The term "carbonylalkyl" represents a carbonyl group attached through an alkyl group having the indicated number of carbon atoms (e.g., acetyl).

[0061] The term "alkylcarbonylalkyl" represents an alkylcarbonyl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., propan-2-one and 4,4-dimethyl-pentan-2-one).

15 [0062] The term "arylcarbonylalkyl" represents a arylcarbonyl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., 1-phenylpropan-1-one and 1-(3-chloro-phenyl)-2-methyl-butan-1-one).

[0063] The term "hetarylcarbonylalkyl" represents a hetarylcarbonyl group as defined above attached through an alkyl group having the indicated number of carbon atoms (e.g., 1-pyridin-2-yl-propan-1-one and 1-(1-*H*-imidazol-2-yl)-propan-1-one).

20 [0064] The term "arylalkylcarbonyl" represents an arylalkyl group as defined above having the indicated number of carbon atoms attached through a carbonyl group (e.g., phenylpropylcarbonyl and phenylethylcarbonyl).

25 [0065] The term "hetarylalkylcarbonyl" represents a hetarylalkyl group as defined above wherein the alkyl group is in turn attached through a carbonyl (e.g., imidazolypentylcarbonyl).

[0066] The term "alkylcarbonylamino" represents an "alkylcarbonyl" group as defined above wherein the carbonyl is in turn attached through the nitrogen atom of an amino group (e.g., methylcarbonylamino, cyclopentylcarbonyl-aminomethyl, and methylcarbonylaminophenyl). The nitrogen atom may itself be substituted with an alkyl or aryl group.

30 [0067] The term "alkylcarbonylaminoalkyl" represents an "alkylcarbonylamino" group attached through an alkyl group having the indicated number of carbon atoms (e.g. N-propylacetamide and N-butyl-propionamide).

[0068] The term "arylalkylcarbonylamino" represents an "arylalkylcarbonyl" group as defined above attached through an amino group (e.g., phenylacetamide and 3-phenyl-propionamide).

[0069] The term "arylalkylcarbonylaminoalkyl" represents an "arylalkylcarbonylamino" group attached through an alkyl group having the indicated number of carbon atoms (e.g., N-ethyl-phenylacetamide and N-butyl-3-phenyl-propionamide).

5 [0070] The term "arylcarbonylamino" represents an "arylcarbonyl" group as defined above attached through an amino group (e.g., benzamide and naphthalene-1-carboxylic acid amide).

[0071] The term "arylcarbonylaminoalkyl" represents an "arylcarbonylamino" group attached through an alkyl group having the indicated number of carbon atoms (e.g., N-propyl-benzamide and N-Butyl-naphthalene-1-carboxylic acid amide).

10 [0072] The term "alkylcarboxy" represents an alkylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen bridge (e.g., heptylcarboxy, cyclopropyl-carboxy, and 3-pentenylcarboxy).

[0073] The term "arylcarboxy" represents an arylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen bridge (e.g., benzoic acid).

15 [0074] The term "alkylcarboxyalkyl" represents an alkylcarboxy group as defined above wherein the oxygen is attached via an alkyl bridge (e.g., heptylcarboxymethyl, propylcarboxy *tert*-butyl, and 3-pentylcarboxyethyl).

20 [0075] The term "arylalkylcarboxy" represents an arylalkylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen bridge (e.g., benzylcarboxy and phenylpropylcarboxy).

[0076] The term "arylalkylcarboxyalkyl" represents an arylalkylcarboxy group as defined above wherein the carboxy group is in turn attached through an alkyl group having the indicated number of carbon atoms (e.g., benzylcarboxymethyl and phenylpropylcarboxypropyl).

25 [0077] The term "hetarylcarboxy" represents a hetarylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen bridge (e.g., pyridine-2-carboxylic acid).

[0078] The term "hetarylalkylcarboxy" represents a hetarylalkylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen bridge (e.g., (1-*H*-imidazol-2-yl)-acetic acid and 3-pyrimidin-2-yl-propionic acid).

30 [0079] The term "alkylSO_m" represents an alkyl group having the number of indicated carbon atoms, wherein the alkyl group is in turn attached through a sulphur bridge wherein the sulphur is substituted with m oxygen atoms (e.g., ethylsulfonyl and ethylsulfinyl).

35 [0080] The term "arylSO_m" represents an aryl group as defined above, wherein the aryl group is in turn attached through a sulphur bridge wherein the sulphur is substituted with m oxygen atoms (e.g., phenylsulfinyl and naphthyl-2-sulfonyl).

[0081] The term "hetarylSO_m" represents a hetaryl group as defined above, wherein the hetaryl group is in turn attached through a sulphur bridge wherein the sulphur is substituted with m oxygen atoms (e.g., thiazol-2-sulfinyl and pyridine-2-sulfonyl).

5 [0082] Certain of the above defined terms may occur more than once in the structural formulae, and upon such occurrence each term shall be defined independently of the other.

[0083] The term "optionally substituted" as used herein means that the groups in question are either unsubstituted or substituted with one or more of the substituents specified. When the groups in question are substituted with more than one substituent, the substituents may be the same or different.

10 [0084] The term "treatment" or "treating" is defined as the management and care of a patient for the purpose of combating or alleviating the disease, condition, or disorder, and the term includes the administration of the active compound to prevent or delay the onset of the symptoms or complications; alleviating (both temporary and permanent) the symptoms or complications; and/or eliminating the disease, condition, or disorder. Thus, "treatment" or "treating"

15 includes prevention and/or prophylaxis of the disease, condition, or disorder.


[0085] The term "pharmaceutically acceptable" is defined as being suitable for administration to humans without adverse events.

20 [0086] The term "prodrug" is defined as a chemically modified form of the active drug, said prodrug being administered to the patient and subsequently being converted to the active drug. Techniques for development of prodrugs are well known in the art.

DETAILED DESCRIPTION OF THE INVENTION

[0087] Thus in an embodiment, the present invention provides for the novel use of a substituted amide, a prodrug thereof, or a salt thereof with a pharmaceutically acceptable acid or

25 base, or any optical isomer or mixture of optical isomers, including a racemic mixture or any tautomeric forms, wherein the substituted amide or a prodrug thereof is of formula I:




[0088] wherein:

[0089] X is selected from CR⁵ and N;

[0090] R¹ is selected from H and C₁-C₆alkyl-R⁶, wherein the alkyl group is substituted with 0-3 R⁷;

5

[0091] R² is selected from hydrogen, halo, C₁-C₆alkyl, and -C(=O)R¹³;



[0092] alternatively, R¹ and R² are, independently,

10 [0093] Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulphur;

15 [0094] Ring A is substituted with 0-3 groups selected from C₁-C₈alkyl, halo, hydroxy, oxo, cyano, C₁-C₆alkoxy, C₁-C₆alkoxyC₁-C₆alkylene, and C₁-C₆alkylcarbonyl, wherein each alkyl/alkylene group is substituted with 0-3 R¹⁴;

[0095] R⁵ is selected from hydrogen, C₁-C₆alkyl, -C(=O)R¹³, and cyano;

20 [0096] R⁶ is selected from cyano, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹, -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₁₀cycloalkyl, -N(R¹⁸)C(=O)-3-10 membered hetcycloalkyl, -N(R¹⁸)C(=O)-aryl, -N(R¹⁸)C(=O)-hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R¹⁶;

25

[0097] R⁷ is selected from halo, hydroxy, oxo, cyano, and C₁-C₈alkyl;

30 [0098] R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from hydrogen, C₁-C₈alkyl, F, trihalomethyl, trihalomethoxy, hydroxy, and C₁-C₆alkoxy, wherein the C₁-C₈alkyl and C₁-C₆alkoxy are substituted with 0-3 R¹⁷;

[0099] alternatively, R⁸ and R⁹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional

carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

5

[00100] alternatively, R¹⁰ and R¹¹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

10

[00101] alternatively, R⁸ and R¹⁰ together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms, 1-4 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy, and hetarylC₁-C₆alkyloxy;

15

[00102] R¹² is selected from H, OH, NR¹⁸R¹⁹, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, and -C(=NR¹⁵)NR¹⁶; wherein the cycloalkyl and hetcycloalkyl groups are substituted with 0-3 R¹⁷;

20

[00103] R¹³ is selected from OH, C₁-C₈alkyl, C₁-C₈alkyloxy, C₁-C₈alkyloxyC₁-C₄alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and NR¹⁸R¹⁹;

25

[00104] R¹⁴ is selected from halo, hydroxy, oxo, and cyano;

[00105] R¹⁵ and R¹⁶ are independently selected from H, C₁-C₈alkyl, 3-10 membered cycloalkyl, halo, OH, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-3 R²⁰;

30

[00106] R¹⁷ is selected from halo, OH, oxo, nitro, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, NR¹⁸R¹⁹, C₁-C₈alkyl, C₁-C₆alkyloxy, and aryloxy;

35

[00107] R^{18} and R^{19} are independently selected from H, C_1 - C_8 alkyl, C_1 - C_8 alkyloxy, aryl, hetaryl, aryl C_1 - C_6 alkylene, and hetaryl C_1 - C_6 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-3 R^{20} ;

- 5 [00108] alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-3 C_1 - C_8 alkyl, aryl, hetaryl, aryl C_1 - C_6 alkylene, hetaryl C_1 - C_6 alkylene, hydroxy, oxo, C_1 - C_6 alkyloxy, aryl C_1 - C_6 alkyloxy, 10 hetaryl C_1 - C_6 alkyloxy, C_1 - C_6 alkyloxy C_1 - C_6 alkyl, C_1 - C_6 alkylcarbonyl, arylcarbonyl, hetarylcarbonyl, aryl C_1 - C_6 alkylcarbonyl, hetaryl C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarboxy, arylcarboxy, hetarylcarboxy, aryl C_1 - C_6 alkyl-carboxy, and hetaryl C_1 - C_6 alkylcarboxy;

- [00109] R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_6 alkyl, C_1 - C_6 alkyloxy, 15 $NR^{21}R^{22}$, methylenedioxy, dihalomethylenedioxy, trihalomethyl, and trihalomethyloxy;

[00110] R^{21} and R^{22} are independently selected from H, C_1 - C_8 alkyl, and aryl C_1 - C_6 alkyl;

- [00111] R^{23} is selected from H and C_1 - C_6 alkyl; 20

[00112] n is selected from 0, 1, and 2;


[00113] Y is selected from O and S;

- 25 [00114] or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

- [00115] In another embodiment, the present invention provides the novel use of compounds 30 of formula I, wherein:

[00116] R^1 is selected from H and C_1 - C_4 alkyl- R^6 , wherein the alkyl group is substituted with 0-1 R^7 ;

- 35 [00117] R^2 is selected from hydrogen, C_1 - C_6 alkyl, and $-C(=O)R^{13}$;



[00118] alternatively, R¹ and R² are, independently,

[00119] Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen and 7-10 carbon atoms;

[00120] Ring A is substituted with 0-3 groups selected from C₁-C₄alkyl, halo, hydroxy, and C₁-C₆alkyloxy;

[00121] R⁵ is selected from hydrogen and C₁-C₄alkyl;

[00122] R⁶ is selected from cyano, C₃-C₆cycloalkyl, 3-6 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹, -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₆cycloalkyl, -N(R¹⁸)C(=O)-3-6 membered hetcycloalkyl, -N(R¹⁸)C(=O)-aryl, -N(R¹⁸)C(=O)-hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R¹⁶;

[00123] R⁷ is selected from halo and C₁-C₄alkyl;

[00124] R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from hydrogen and C₁-C₄alkyl;

[00125] alternatively, R⁸ and R¹⁰ together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms and 1-4 additional carbon atoms, wherein this ring is substituted with 0-1 groups selected from halo, trihalomethyl, hydroxyl, and C₁-C₆alkyl;

[00126] R¹² is selected from H, OH, and NR¹⁸R¹⁹;

[00127] R¹³ is selected from OH, C₁-C₄alkyl, C₁-C₄alkyloxy, C₁-C₄alkyloxyC₁-C₄alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and NR¹⁸R¹⁹;

[00128] R^{15} and R^{16} are independently selected from H, C_1 - C_4 alkyl, 3-6 membered cycloalkyl, halo, OH, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-1 R^{20} ;

5 **[00129]** R^{18} and R^{19} are independently selected from H, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, aryl, hetaryl, aryl C_1 - C_4 alkylene, and hetaryl C_1 - C_4 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-1 R^{20} ;

10 **[00130]** alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-5 carbon atoms, and 0-1 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-1 C_1 - C_4 alkyl, aryl, hetaryl, aryl C_1 - C_4 alkylene, hetaryl C_1 - C_4 alkylene, hydroxy, and C_1 - C_4 alkyloxy;

15 **[00131]** R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, $NR^{21}R^{22}$, trihalomethyl, and trihalomethyloxy;

[00132] R^{21} and R^{22} are independently selected from H, C_1 - C_4 alkyl, and aryl C_1 - C_4 alkyl;


20 **[00133]** R^{23} is selected from H and C_1 - C_6 alkyl;

[00134] n is selected from 0, 1, and 2; and,

[00135] Y is selected from O and S.

25


[00136] In another embodiment, the present invention provides the novel use of compounds wherein the substituted amide or prodrug thereof is of formula IA:



IA.


30

[00137] In another embodiment, the present invention provides the novel use of compounds wherein the substituted amide or prodrug thereof is of formula IB:




IB.

[00138] In another embodiment, the present invention provides the novel use of compounds wherein the substituted amide or prodrug thereof is of formula IC:



IC.


[00139] In another embodiment, the present invention provides the novel use of compounds wherein the substituted amide or prodrug thereof is of formula ID:



ID.

[00140] In another embodiment, the present invention provides the novel use of compounds of formula I, wherein:

[00141] Ring A is selected from:




5

[00142] Ring A is substituted with 0-2 R^{24} , and,

[00143] R^{24} is selected from C_1 - C_8 alkyl, halo, hydroxy, oxo, cyano, and C_1 - C_6 alkyloxy.

10

[00144] In another embodiment, the present invention provides the novel use of compounds of formula I, wherein:



15

[00145] Ring A is

[00146] In another embodiment, the present invention provides the novel use of compounds of formula I, wherein the substituted amide or a prodrug thereof is of the selected from the group:

20

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester;

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid

25

[00147] or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

[00148] In another embodiment, the present invention provides for the novel preparation of a pharmaceutical composition for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

5

[00149] In another embodiment, the present invention provides for the novel preparation of a pharmaceutical composition, wherein: the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.

10 **[00150]** In another embodiment, the present invention provides for the novel preparation of a pharmaceutical composition, wherein: the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), the progression from IGT to type 2 diabetes, the progression of the metabolic syndrome into type 2 diabetes,
15 diabetic late complications, neurodegenerative and psychiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

[00151] In another embodiment, the present invention provides for the novel preparation of a pharmaceutical composition, wherein: the pharmaceutical composition is suitable for a route
20 of administration selected from oral, nasal, buccal, transdermal, pulmonal, and parenteral.

[00152] In another embodiment, the present invention provides a novel method for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial, the method comprising administering to a subject in need
25 thereof an effective amount of a compound of the present invention.


[00153] In another embodiment, the present invention provides a novel method wherein the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.

30 **[00154]** In another embodiment, the present invention provides a novel method wherein the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), progression from IGT to type 2 diabetes, progression of metabolic syndrome into type 2 diabetes, diabetic late complications, neurodegenerative and psy-

chiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

[00155] In another embodiment, the present invention provides a novel method wherein the administering is via a route selected from oral, nasal, buccal, transdermal, pulmonal, and parenteral.

[00156] In another embodiment, the present invention provides for a novel compound of the formula I:




[00157] wherein:

[00158] X is selected from CR⁵ and N;

[00159] R¹ is selected from H and C₁-C₆alkyl-R⁶, wherein the alkyl group is substituted with 0-3 R⁷;

[00160] R² is selected from hydrogen, halo, C₁-C₆alkyl, and -C(=O)R¹³;

[00161] alternatively, R¹ and R² are, independently,



[00162] Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulphur;

[00163] Ring A is substituted with 0-3 groups selected from C₁-C₈alkyl, halo, hydroxy, oxo, cyano, C₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkylene, and C₁-C₆alkylcarbonyl, wherein each alkyl/alkylene group is substituted with 0-3 R¹⁴;

5 **[00164]** R⁵ is selected from hydrogen, C₁-C₆alkyl, -C(=O)R¹³, and cyano;

[00165] R⁶ is selected from cyano, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹, -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₁₀cycloalkyl, -N(R¹⁸)C(=O)-3-10 membered hetcycloalkyl, -N(R¹⁸)C(=O)-aryl, -N(R¹⁸)C(=O)-hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R¹⁶;

10

[00166] R⁷ is selected from halo, hydroxy, oxo, cyano, and C₁-C₈alkyl;

15 **[00167]** R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from hydrogen, C₁-C₈alkyl, F, trihalomethyl, trihalomethoxy, hydroxy, and C₁-C₆alkyloxy, wherein the C₁-C₈alkyl and C₁-C₆alkyloxy are substituted with 0-3 R¹⁷;

[00168] alternatively, R⁸ and R⁹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

20

[00169] alternatively, R¹⁰ and R¹¹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

25

30

[00170] alternatively, R⁸ and R¹⁰ together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms, 1-4 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sul-

35

phur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy, and hetarylC₁-C₆alkyloxy;

- 5 **[00171]** R¹² is selected from H, OH, NR¹⁸R¹⁹, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, and -C(=NR¹⁵)NR¹⁶; wherein the cycloalkyl and hetcycloalkyl groups are substituted with 0-3 R¹⁷;

- 10 **[00172]** R¹³ is selected from OH, C₁-C₈alkyl, C₁-C₈alkyloxy, C₁-C₈alkyloxyC₁-C₄alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and NR¹⁸R¹⁹;

[00173] R¹⁴ is selected from halo, hydroxy, oxo, and cyano;

- 15 **[00174]** R¹⁵ and R¹⁶ are independently selected from H, C₁-C₈alkyl, 3-10 membered cycloalkyl, halo, OH, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-3 R²⁰;

- 20 **[00175]** R¹⁷ is selected from halo, OH, oxo, nitro, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, NR¹⁸R¹⁹, C₁-C₈alkyl, C₁-C₆alkyloxy, and aryloxy;

[00176] R¹⁸ and R¹⁹ are independently selected from H, C₁-C₈alkyl, C₁-C₈alkyloxy, aryl, hetaryl, arylC₁-C₆alkylene, and hetarylC₁-C₆alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-3 R²⁰;

- 25 **[00177]** alternatively, R¹⁸ and R¹⁹, together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-3 C₁-C₈alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, arylC₁-C₆alkyloxy, 30 hetarylC₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkyl, C₁-C₆alkylcarbonyl, arylcarbonyl, hetarylcarbonyl, arylC₁-C₆alkylcarbonyl, hetarylC₁-C₆alkylcarbonyl, C₁-C₆alkylcarboxy, arylcarboxy, hetarylcarboxy, arylC₁-C₆alkyl-carboxy, and hetarylC₁-C₆alkylcarboxy;

- 35 **[00178]** R²⁰ is selected from H, OH, oxo, halo, cyano, nitro, C₁-C₆alkyl, C₁-C₆alkyloxy, NR²¹R²², methylenedioxy, dihalomethylenedioxy, trihalomethyl, and trihalomethyloxy;

[00179] R^{21} and R^{22} are independently selected from H, C_1 - C_8 alkyl, and aryl C_1 - C_6 alkyl;

[00180] R^{23} is selected from H and C_1 - C_6 alkyl;

5

[00181] n is selected from 0, 1, and 2;

[00182] Y is selected from O and S;

10 [00183] or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

[00184] In another embodiment, the present invention provides the novel compounds of formula I, wherein:


15

[00185] R^1 is selected from H and C_1 - C_4 alkyl- R^6 , wherein the alkyl group is substituted with 0-1 R^7 ;

[00186] R^2 is selected from hydrogen, C_1 - C_6 alkyl, and $-C(=O)R^{13}$;

20

[00187] alternatively, R^1 and R^2 are, independently,



[00188] Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen and 7-10 carbon atoms;

25

[00189] Ring A is substituted with 0-3 groups selected from C_1 - C_4 alkyl, halo, hydroxy, and C_1 - C_6 alkyloxy;

[00190] R^5 is selected from hydrogen and C_1 - C_4 alkyl;

30

[00191] R^6 is selected from cyano, C_3 - C_6 cycloalkyl, 3-6 membered hetcycloalkyl, aryl, hetaryl, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $-S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, $-N(R^{23})C(=Y)NR^{18}R^{19}$, $-C(=NR^{15})NR^{15}$, $-N(R^{18})C(=O)R^{13}$, $-N(R^{18})C(=O)-C_3$ - C_6 cycloalkyl, $-N(R^{18})C(=O)$ -3-6 mem-

bered hetcycloalkyl, $-N(R^{18})C(=O)$ -aryl, $-N(R^{18})C(=O)$ -hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R^{16} ;

[00192] R^7 is selected from halo and C_1 - C_4 alkyl;

5

[00193] R^8 , R^9 , R^{10} and R^{11} are each independently selected from hydrogen and C_1 - C_4 alkyl;

[00194] alternatively, R^8 and R^{10} together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms and 1-4 additional carbon atoms, wherein this ring is substituted with 0-1 groups selected from halo, trihalomethyl, hydroxyl, and C_1 - C_6 alkyl;

10

[00195] R^{12} is selected from H, OH, and $NR^{18}R^{19}$;

15

[00196] R^{13} is selected from OH, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, C_1 - C_4 alkyloxy C_1 - C_4 alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and $NR^{18}R^{19}$;

[00197] R^{15} and R^{16} are independently selected from H, C_1 - C_4 alkyl, 3-6 membered cycloalkyl, halo, OH, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-1 R^{20} ;

20

[00198] R^{18} and R^{19} are independently selected from H, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, aryl, hetaryl, aryl C_1 - C_4 alkylene, and hetaryl C_1 - C_4 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-1 R^{20} ;

25

[00199] alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-5 carbon atoms, and 0-1 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-1 C_1 - C_4 alkyl, aryl, hetaryl, aryl C_1 - C_4 alkylene, hetaryl C_1 - C_4 alkylene, hydroxy, and C_1 - C_4 alkyloxy;

30

[00200] R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, $NR^{21}R^{22}$, trihalomethyl, and trihalomethyloxy;

35

[00201] R^{21} and R^{22} are independently selected from H, C_1 - C_4 alkyl, and aryl C_1 - C_4 alkyl;

[00202] R^{23} is selected from H and C_1 - C_6 alkyl;


[00203] n is selected from 0, 1, and 2; and,

5

[00204] Y is selected from O and S.

[00205] In another embodiment, the present invention provides the novel compounds of formula IA:


10



IA.

[00206] In another embodiment, the present invention provides the novel compounds of formula IB:


15



IB.

20


[00207] In another embodiment, the present invention provides the novel compounds of formula IC:



IC.

5

[00208] In another embodiment, the present invention provides the novel compounds of formula ID:




ID.

10

[00209] In another embodiment, the present invention provides the novel compounds of formula I, wherein:

15


[00210] Ring A is selected from:



[00211] Ring A is substituted with 0-2 R²⁴; and,

[00212] R^{24} is selected from C_1 - C_8 alkyl, halo, hydroxy, oxo, cyano, and C_1 - C_6 alkyloxy.

- 5 [00213] In another embodiment, the present invention provides the novel compounds of formula I, wherein:



[00214] Ring A is

10

[00215] In another embodiment, the present invention provides the novel compounds of formula I, selected from the group:

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester;

- 15 3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid;

[00216] or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

- 20 [00217] In another embodiment, the present invention provides a novel compound, which is an agent useful for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

- 25 [00218] In another embodiment, the present invention provides a novel method wherein the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.

- 30 [00219] In another embodiment, the present invention provides a novel method wherein the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), progression from IGT to type 2 diabetes, progression of metabolic syndrome into type 2 diabetes, diabetic late complications, neurodegenerative and psychiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

[00220] In another embodiment, the present invention provides a novel pharmaceutical composition comprising, as an active ingredient, at least one compound according to the present invention together with one or more pharmaceutically acceptable carriers or excipients.

[00221] In another embodiment, the present invention provides a novel pharmaceutical composition, which is suitable for oral, nasal, buccal, transdermal, pulmonary, or parenteral administration.

[00222] The compounds of the present invention may have asymmetric centers and may occur as racemates, racemic mixtures, and as individual enantiomers or diastereoisomers, with all isomeric forms being included in the present invention as well as mixtures thereof.

[00223] The present invention also encompasses pharmaceutically acceptable salts of the present compounds. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable base addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, and nitric acids. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pantoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, sulphates, nitrates, phosphates, perchlorates, borates, acetates, benzoates, hydroxynaphthoates, glycerophosphates, and ketoglutarates. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci., 66, 2 (1977), which is incorporated herein by reference. Examples of metal salts include lithium, sodium, potassium, barium, calcium, magnesium, zinc, and calcium salts. Examples of amines and organic amines include ammonium, methylamine, dimethylamine, trimethylamine, ethylamine, diethylamine, propylamine, butylamine, tetramethylamine, ethanolamine, diethanolamine, triethanolamine, meglumine, ethylenediamine, choline, N,N'-dibenzylethylenediamine, N-benzylphenylethylamine, N-methyl-D-glucamine, and guanidine. Examples of cationic amino acids include lysine, arginine, and histidine.

[00224] Further, some of the compounds of the present invention may form solvates with water or common organic solvents. Such solvates are encompassed within the scope of the invention.

[00225] The pharmaceutically acceptable salts are prepared by reacting a compound of the present invention with 1 to 4 equivalents of a base such as sodium hydroxide, sodium methoxide, sodium hydride, potassium *tert*-butoxide, calcium hydroxide, and magnesium hydroxide, in solvents such as ether, THF, methanol, *tert*-butanol, dioxane, and isopropanol, ethanol. Mixtures of solvents may be used. Organic bases such as lysine, arginine, diethanolamine, choline, guandine and their derivatives etc. may also be used. Alternatively, acid addition salts wherever applicable are prepared by treatment with acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, p-toluenesulphonic acid, methanesulfonic acid, acetic acid, citric acid, maleic acid, salicylic acid, hydroxynaphthoic acid, ascorbic acid, palmitic acid, succinic acid, benzoic acid, benzenesulfonic acid, and tartaric acid in solvents such as ethyl acetate, ether, alcohols, acetone, THF, and dioxane. Mixture of solvents may also be used.

[00226] The stereoisomers of the compounds forming part of this invention may be prepared by using reactants in their single enantiomeric form in the process wherever possible or by conducting the reaction in the presence of reagents or catalysts in their single enantiomer form or by resolving the mixture of stereoisomers by conventional methods. Some of the preferred methods include use of microbial resolution, enzymatic resolution, resolving the diastereomeric salts formed with chiral acids such as mandelic acid, camphorsulfonic acid, tartaric acid, and lactic acid, wherever applicable or chiral bases such as brucine, (*R*)- or (*S*)-phenylethylamine, cinchona alkaloids and their derivatives. Commonly used methods are compiled by Jaques et al. in "Enantiomers, Racemates and Resolution" (Wiley Interscience, 1981). More specifically the compound of the present invention may be converted to a 1:1 mixture of diastereomeric amides by treating with chiral amines, aminoacids, aminoalcohols derived from aminoacids; conventional reaction conditions may be employed to convert acid into an amide; the diastereomers may be separated either by fractional crystallization or chromatography and the stereoisomers of compound of formula I may be prepared by hydrolysing the pure diastereomeric amide.

[00227] Various polymorphs of the compounds forming part of this invention may be prepared by crystallization of said compounds under different conditions. For example, using different solvents commonly used or their mixtures for recrystallization; crystallizations at different temperatures; various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Polymorphs may also be obtained by heating or melting the com-

pound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, ir spectroscopy, differential scanning calorimetry, powder X-ray diffraction or such other techniques.

[00228] The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming active pharmacological substances. In general, such prodrugs will be functional derivatives of the present compounds, which are readily convertible *in vivo* into the required compound of the present invention. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

[00229] It is a well known problem in drug discovery that compounds, such as enzyme inhibitors, may be very potent and selective in biochemical assays, yet be inactive *in vivo*. This lack of so-called bioavailability may be ascribed to a number of different factors such as lack of or poor absorption in the gut, first pass metabolism in the liver and/or poor uptake in cells. Although the factors determining bioavailability are not completely understood, there are many examples in the scientific literature - well known to those skilled in the art - of how to modify compounds, which are potent and selective in biochemical assays but show low or no activity *in vivo*, into drugs that are biologically active.

[00230] It is within the scope of the invention to modify the compounds of the present invention, termed the 'original compound', by attaching chemical groups that will improve the bioavailability of said compounds in such a way that the uptake in cells or mammals is facilitated.

[00231] Examples of said modifications, which are not intended in any way to limit the scope of the invention, include changing of one or more carboxy groups to esters (for instance methyl esters, ethyl esters, *tert*-butyl, acetoxymethyl, pivaloyloxymethyl esters or other acyloxymethyl esters). Compounds of the invention, original compounds, such modified by attaching chemical groups are termed 'modified compounds'.

[00232] The invention also encompasses active metabolites of the present compounds.

[00233] The compounds according to the invention alter, and more specifically, reduce the level of active intracellular glucocorticoid and are accordingly useful for the treatment of conditions, disorders, and diseases in which such a modulation or reduction is beneficial.

[00234] Accordingly, the present compounds may be applicable for the treatment of the metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), Latent Autoimmune Diabetes in the Adult (LADA), type 1 diabetes, diabetic late complications including cardiovascular

diseases, cardiovascular disorders, disorders of lipid metabolism, neurodegenerative and psychiatric disorders, dysregulation of intraocular pressure including glaucoma, immune disorders, inappropriate immune responses, musculo-skeletal disorders, gastrointestinal disorders, polycystic ovarie syndrome (PCOS), reduced hair growth or other diseases, disorders or conditions that are influenced by intracellular glucocorticoid levels, adverse effects of increased blood levels of active endogenous or exogenous glucocorticoid, and any combination thereof, adverse effects of increased plasma levels of endogenous active glucocorticoid, Cushing's disease, Cushing's syndrome, adverse effects of glucocorticoid receptor agonist treatment of autoimmune diseases, adverse effects of glucocorticoid receptor agonist treatment of inflammatory diseases, adverse effects of glucocorticoid receptor agonist treatment of diseases with an inflammatory component, adverse effects of glucocorticoid receptor agonist treatment as a part of cancer chemotherapy, adverse effects of glucocorticoid receptor agonist treatment for surgical/post-surgical or other trauma, adverse effects of glucocorticoid receptor agonist therapy in the context of organ or tissue transplantation or adverse effects of glucocorticoid receptor agonist treatment in other diseases, disorders or conditions where glucocorticoid receptor agonists provide clinically beneficial effects. Also the present compounds may be applicable for thr treatment of visceral fat accumulation and insulin resistance in HAART (highly active antiretroviral treatment)-treated patients.

[00235] More specifically the present compounds may be applicable for the treatment of the metabolic syndrome, type 2 diabetes, diabetes as a consequence of obesity, insulin resistance, hyperglycemia, prandial hyperglycemia, hyperinsulinemia, inappropriately low insulin secretion, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), increased hepatic glucose production, type 1 diabetes, LADA, pediatric diabetes, dyslipidemia, diabetic dyslipidemia, hyperlipidemia, hypertriglyceridemia, hyperlipoproteinemia, hypercholesterolemia, decreased HDL cholesterol, impaired LDL/HDL ratio, other disorders of lipid metabolism, obesity, visceral obesity, obesity as a consequence of diabetes, increased food intake, hypertension, diabetic late complications, micro-/macroalbuminuria, nephropathy, retinopathy, neuropathy, diabetic ulcers, cardiovascular diseases, arteriosclerosis, atherosclerosis, coronary artery disease, cardiac hypertrophy, myocardial ischemia, heart insufficiency, congestional heart failure, stroke, myocardial infarction, arrhythmia, decreased blood flow, erectile dysfunction (male or female), myopathy, loss of muscle tissue, muscle wasting, muscle catabolism, osteoporosis, decreased linear growth, neurodegenerative and psychiatric disorders, Alzheimers disease, neuronal death, impaired cognitive function, depression, anxiety, eating disorders, appetite regulation, migraine, epilepsy, addiction to chemical substances, disorders of intraocular pressure, glaucoma, polycystic ovary syndrome (PCOS),

inappropriate immune responses, inappropriate T helper-1/T helper-2 polarisation, bacterial infections, mycobacterial infections, fungal infections, viral infections, parasitic infestations, suboptimal responses to immunizations, immune dysfunction, partial or complete baldness, or other diseases, disorders or conditions that are influenced by intracellular glucocorticoid levels and any combination thereof, adverse effects of glucocorticoid receptor agonist treatment of allergic-inflammatory diseases such as asthma and atopic dermatitis, adverse effects of glucocorticoid receptor agonist treatment of disorders of the respiratory system e.g., asthma, cystic fibrosis, emphysema, bronchitis, hypersensitivity, pneumonitis, eosinophilic pneumonias, pulmonary fibrosis, adverse effects of glucocorticoid receptor agonist treatment of inflammatory bowel disease such as Crohn's disease and ulcerative colitis; adverse effects of glucocorticoid receptor agonist treatment of disorders of the immune system, connective tissue and joints e.g., reactive arthritis, rheumatoid arthritis, Sjögren's syndrome, systemic lupus erythematosus, lupus nephritis, Henoch-Schönlein purpura, Wegener's granulomatosis, temporal arteritis, systemic sclerosis, vasculitis, sarcoidosis, dermatomyositis-polymyositis, pemphigus vulgaris; adverse effects of glucocorticoid receptor agonist treatment of endocrinological diseases such as hyperthyroidism, hypoadosteronism, hypopituitarism; adverse effects of glucocorticoid receptor agonist treatment of hematological diseases e.g., hemolytic anemia, thrombocytopenia, paroxysmal nocturnal hemoglobinuria; adverse effects of glucocorticoid receptor agonist treatment of cancer such as spinal cord diseases, neoplastic compression of the spinal cord, brain tumours, acute lymphoblastic leukemia, Hodgkin's disease, chemotherapy-induced nausea, adverse effects of glucocorticoid receptor agonist treatment of diseases of muscle and at the neuro-muscular joint e.g., myasthenia gravis and hereditary myopathies (e.g., Duchenne muscular dystrophy), adverse effects of glucocorticoid receptor agonist treatment in the context of surgery & transplantation e.g., trauma, post-surgical stress, surgical stress, renal transplantation, liver transplantation, lung transplantation, pancreatic islet transplantation, blood stem cell transplantation, bone marrow transplantation, heart transplantation, adrenal gland transplantation, tracheal transplantation, intestinal transplantation, corneal transplantation, skin grafting, keratoplasty, lens implantation and other procedures where immunosuppression with glucocorticoid receptor agonists is beneficial; adverse effects of glucocorticoid receptor agonist treatment of brain abscess, nausea/vomiting, infections, hypercalcemia, adrenal hyperplasia, autoimmune hepatitis, spinal cord diseases, saccular aneurysms or adverse effects to glucocorticoid receptor agonist treatment in other diseases, disorders and conditions where glucocorticoid receptor agonists provide clinically beneficial effects.

[00236] Accordingly, in a further aspect the invention relates to a compound according to the invention for use as a pharmaceutical composition.

[00237] The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound according to the invention together with one or more pharmaceutically acceptable carriers or diluents.

[00238] The pharmaceutical composition is preferably in unit dosage form, comprising from about 0.05 mg/day to about 2000 mg/day, preferably from about 0.1 mg/day to about 1000 mg/day, and more preferably from about 0.5 mg/day to about 500 mg/day of a compound according to the invention.

[00239] In another embodiment, the patient is treated with a compound according to the invention for at least about 1 week, for at least about 2 weeks, for at least about 4 weeks, for at least about 2 months or for at least about 4 months.

[00240] In yet another embodiment, the pharmaceutical composition is for oral, nasal, buccal, transdermal, pulmonary or parenteral administration.

[00241] Furthermore, the invention relates to the use of a compound according to the invention for the preparation of a pharmaceutical composition for the treatment of disorders and diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

[00242] The invention also relates to a method for the treatment of disorders and diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial, the method comprising administering to a subject in need thereof an effective amount of a compound according to the invention.

[00243] In a preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the treatment of any diseases and conditions that are influenced by intracellular glucocorticoid levels as mentioned above.

[00244] Thus, in a preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the treatment of conditions and disorders where a decreased level of active intracellular glucocorticoid is desirable, such as the conditions and diseases mentioned above.

[00245] In yet a preferred embodiment of the invention the present compounds are used for the preparation of a medicament for the treatment of metabolic syndrome, insulin resistance, dyslipidemia, hypertension obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), progression from IGT to type 2 diabetes, progression of the metabolic syndrome into type 2 diabetes, diabetic late complications (e.g., cardiovascular diseases, arteriosclerosis, and atherosclerosis), neurodegenerative and psychiatric disorders, and, the adverse effects of glucocorticoid receptor agonist treatment or therapy.

[00246] In another embodiment of the present invention, the route of administration may be any route which effectively transports a compound according to the invention to the appropriate or desired site of action, such as oral, nasal, buccal, transdermal, pulmonal, or parenteral.

5 **[00247]** In still a further aspect of the invention the present compounds are administered in combination with one or more further active substances in any suitable ratios. Such further active substances may e.g., be selected from antiobesity agents, antidiabetics, agents modifying the lipid metabolism, antihypertensive agents, glucocorticoid receptor agonists, agents for the treatment and/or prevention of complications resulting from or associated with diabe-
10 tes and agents for the treatment and/or prevention of complications and disorders resulting from or associated with obesity.

[00248] Thus, in a further aspect of the invention the present compounds may be administered in combination with one or more antiobesity agents or appetite regulating agents.

[00249] Such agents may be selected from the group consisting of CART (cocaine am-
15 phetamine regulated transcript) agonists, NPY (neuropeptide Y) antagonists, MC4 (melanocortin 4) agonists, orexin antagonists, TNF (tumor necrosis factor) agonists, CRF (corticotropin releasing factor) agonists, CRF BP (corticotropin releasing factor binding protein) antagonists, urocortin agonists, $\beta 3$ agonists, MSH (melanocyte-stimulating hormone) agonists, MCH (melanocyte-concentrating hormone) antagonists, CCK (cholecystokinin) agonists, se-
20 rotonin re-uptake inhibitors, serotonin and noradrenaline re-uptake inhibitors, mixed serotonin and noradrenergic compounds, 5HT (serotonin) agonists, bombesin agonists, galanin antagonists, growth hormone, growth hormone releasing compounds, TRH (thyreotropin releasing hormone) agonists, UCP 2 or 3 (uncoupling protein 2 or 3) modulators, leptin ago-
nists, DA agonists (bromocriptin, doprexin), lipase/amylase inhibitors, PPAR (peroxisome
25 proliferator-activated receptor) modulators, RXR (retinoid X receptor) modulators, TR β agonists, AGRP (Agouti related protein) inhibitors, H3 histamine antagonists, opioid antagonists (such as naltrexone), exendin-4, GLP-1 and ciliary neurotrophic factor.

[00250] In one embodiment of the invention the antiobesity agent is leptin; dexamphetamine or amphetamine; fenfluramine or dexfenfluramine; sibutramine; orlistat; mazindol or phen-
30 termine.

[00251] Suitable antidiabetic agents include insulin, insulin analogues and derivatives such as those disclosed in EP 792 290 (Novo Nordisk A/S), e.g., N^{εB29}-tetradecanoyl des (B30) human insulin, EP 214 826 and EP 705 275 (Novo Nordisk A/S), e.g., Asp^{B28} human insulin, US 5,504,188 (Eli Lilly), e.g., Lys^{B28} Pro^{B29} human insulin, EP 368 187 (Aventis), eg Lantus,
35 which are all incorporated herein by reference, GLP-1 (glucagon like peptide-1) and GLP-1

derivatives such as those disclosed in WO 98/08871 to Novo Nordisk A/S, which is incorporated herein by reference as well as orally active hypoglycaemic agents.

[00252] The orally active hypoglycaemic agents preferably comprise sulphonylureas, biguanides, meglitinides, glucosidase inhibitors, glucagon antagonists such as those disclosed in
5 WO 99/01423 to Novo Nordisk A/S and Agouron Pharmaceuticals, Inc., GLP-1 agonists, potassium channel openers such as those disclosed in WO 97/26265 and WO 99/03861 to Novo Nordisk A/S which are incorporated herein by reference, DPP-IV (dipeptidyl peptidase-IV) inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, compounds modifying the lipid metabolism such
10 as antihyperlipidemic agents and antilipidemic agents as PPAR α modulators, PPAR δ modulators, cholesterol absorption inhibitors, HSL (hormone-sensitive lipase) inhibitors and HMG CoA inhibitors (statins), nicotinic acid, fibrates, anion exchangers, compounds lowering food intake, bile acid resins, RXR agonists and agents acting on the ATP-dependent potassium channel of the β -cells.

[00253] In one embodiment, the present compounds are administered in combination with
15 insulin or an insulin analogue or derivative, such as N^{B29}-tetradecanoyl des (B30) human insulin, Asp^{B28} human insulin, Lys^{B28} Pro^{B29} human insulin, Lantus®, or a mix-preparation comprising one or more of these.

[00254] In a further embodiment the present compounds are administered in combination
20 with a sulphonylurea e.g., tolbutamide, glibenclamide, glipizide or glicazide.

[00255] In another embodiment the present compounds are administered in combination with a biguanide e.g., metformin.

[00256] In yet another embodiment the present compounds are administered in combination with a meglitinide e.g., repaglinide or senaglinide.

[00257] In still another embodiment the present compounds are administered in combination
25 with a thiazolidinedione e.g., troglitazone, ciglitazone, pioglitazone, rosiglitazone or compounds disclosed in WO 97/41097 such as 5-[[4-[3-Methyl-4-oxo-3,4-dihydro-2-quinazolinyl]methoxy]phenyl-methyl]thiazolidine-2,4-dione or a pharmaceutically acceptable salt thereof, preferably the potassium salt.

[00258] In yet another embodiment the present compounds may be administered in combination
30 with the insulin sensitizers disclosed in WO 99/19313 such as (-) 3-[4-[2-Phenoxazin-10-yl)ethoxy]phenyl]-2-ethoxypropanoic acid or a pharmaceutically acceptable salts thereof, preferably the arginine salt.

[00259] In a further embodiment the present compounds are administered in combination
35 with an α -glucosidase inhibitor e.g., miglitol or acarbose.

[00260] In another embodiment the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the β -cells e.g., tolbutamide, glibenclamide, glipizide, glicazide or repaglinide.

[00261] Furthermore, the present compounds may be administered in combination with nateglinide.

[00262] In still another embodiment the present compounds are administered in combination with an antihyperlipidemic agent or antilipidemic agent e.g., cholestyramine, colestipol, clofibrate, gemfibrozil, fenofibrate, bezafibrate, tesaglitazar, EML-4156, LY-818, MK-767, atorvastatin, fluvastatin, lovastatin, pravastatin, simvastatin, acipimox, probucol, ezetimibe or dextrothyroxine.

[00263] In a further embodiment the present compounds are administered in combination with more than one of the above-mentioned compounds e.g., in combination with a sulphonylurea and metformin, a sulphonylurea and acarbose, repaglinide and metformin, insulin and a sulphonylurea, insulin and metformin, insulin, insulin and lovastatin, etc.

[00264] Further, the present compounds may be administered in combination with one or more antihypertensive agents. Examples of antihypertensive agents are β -blockers such as alprenolol, atenolol, timolol, pindolol, propranolol, metoprolol, bisoprololfumerate, esmolol, acebutolol, metoprolol, acebutolol, betaxolol, celiprolol, nebivolol, tertatolol, oxprenolol, amosolalul, carvedilol, labetalol, β 2-receptor blockers e.g., S-atenolol, OPC-1085, ACE (angiotensin converting enzyme) inhibitors such as quinapril, lisinopril, enalapril, captopril, benazepril, perindopril, trandolapril, fosinopril, ramipril, cilazapril, delapril, imidapril, moexipril, spirapril, temocapril, zofenopril, S-5590, fasidotril, Hoechst-Marion Roussel: 100240 (EP 00481522), omapatrilat, gemopatrilat and GW-660511, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem, amlodipine, nitrendipine, verapamil, lacidipine, lercanidipine, aranidipine, cilnidipine, clevidipine, azelnidipine, barnidipine, efonodipine, iasidipine, iemildipine, iercanidipine, manidipine, nilvadipine, pranidipine, furnidipine, α -blockers such as doxazosin, urapidil, prazosin, terazosin, bunazosin and OPC-28326, diuretics such as thiazides/sulphonamides (e.g., bendroflumetazide, chlorothalidone, hydrochlorothiazide and clopamide), loop-diuretics (e.g., bumetanide, furosemide and torasemide) and potassium sparing diuretics (e.g., amiloride, spironolactone), endothelin ET-A antagonists such as ABT-546, ambrisentan, atrasentan, SB-234551, CI-1034, S-0139 and YM-598, endothelin antagonists e.g., bosentan and J-104133, renin inhibitors such as aliskiren, vasopressin V1 antagonists e.g., OPC-21268, vasopressin V2 antagonists such as tolvaptan, SR-121463 and OPC-31260, B-type natriuretic peptide agonists e.g., Nesiritide, angiotensin II antagonists such as irbesartan, candesartancilexetil,

losartan, valsartan, telmisartan, eprosartan, candesartan, CL-329167, eprosartan, iosartan, olmesartan, prazosartan, TA-606, and YM-358, 5-HT₂ agonists e.g., fenoldopam and ketanserin, adenosine A₁ antagonists such as naftopidil, N-0861 and FK-352, thromboxane A₂ antagonists such as KT2-962, endopeptidase inhibitors e.g., ecadotril, nitric oxide
5 agonists such as LP-805, dopamine D₁ antagonists e.g., MYD-37, dopamine D₂ agonists such as nalmefene, n-3 fatty acids e.g., omacor, prostacyclin agonists such as treprostinil, beraprost, PGE₁ agonists e.g., ecraprost, Na⁺/K⁺ ATPase modulators e.g., PST-2238, Potassium channel activators e.g., KR-30450, vaccines such as PMD-3117, Indapamides, CGRP-unigene, guanylate cyclase stimulators, hydralazines, methyldopa, docarpamine,
10 moxonidine, CoAprovel, MondoBiotech-811.

[00265] Further reference can be made to Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

[00266] Furthermore, the present compounds may be administered in combination with one or more glucocorticoid receptor agonists. Examples of such glucocorticoid receptor agonists
15 are betametasone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, beclomethasone, butixicort, clobetasol, flunisolide, flucatisone (and analogues), momethasone, triamcinolonacetone, triamcinolonhexacetone GW-685698, NXC-1015, NXC-1020, NXC-1021, NS-126, P-4112, P-4114, RU-24858 and T-25 series.

[00267] It should be understood that any suitable combination of the compounds according
20 to the invention with one or more of the above-mentioned compounds and optionally one or more further pharmacologically active substances are considered to be within the scope of the present invention.

PHARMACEUTICAL COMPOSITIONS

[00268] The compounds of the present invention may be administered alone or in combination
25 with pharmaceutically acceptable carriers or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co.,
30 Easton, PA, 1995.

[00269] The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as the oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) route, the oral route be-

ing preferred. It will be appreciated that the preferred route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient chosen.

[00270] Pharmaceutical compositions for oral administration include solid dosage forms such as hard or soft capsules, tablets, troches, dragees, pills, lozenges, powders and granules. Where appropriate, they can be prepared with coatings such as enteric coatings or they can be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well-known in the art.

[00271] Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

[00272] Pharmaceutical compositions for parenteral administration include sterile aqueous and non-aqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Depot injectable formulations are also contemplated as being within the scope of the present invention.

[00273] Other suitable administration forms include suppositories, sprays, ointments, crèmes, gels, inhalants, dermal patches, implants etc.

[00274] A typical oral dosage is in the range of from about 0.001 to about 100 mg/kg body weight per day, preferably from about 0.01 to about 50 mg/kg body weight per day, and more preferred from about 0.05 to about 10 mg/kg body weight per day administered in one or more dosages such as 1 to 3 dosages. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

[00275] The formulations may conveniently be presented in unit dosage form by methods known to those skilled in the art. A typical unit dosage form for oral administration one or more times per day such as 1 to 3 times per day may contain from 0.05 to about 2000 mg, e.g., from about 0.1 to about 1000 mg, from about 0.5 mg to about 500 mg., from about 1 mg to about 200 mg, e.g., about 100 mg.

[00276] For parenteral routes, such as intravenous, intrathecal, intramuscular and similar administration, typically doses are in the order of about half the dose employed for oral administration.

[00277] The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. Examples are an acid addition salt of a compound having the utility of a free base and a base addition salt of a compound having the utility of a

free acid. The term "pharmaceutically acceptable salts" refers to non-toxic salts of the compounds for use according to the present invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid or by reacting the acid with a suitable organic or inorganic base. When a compound for use according to the present invention, contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of the compound with a chemical equivalent of a pharmaceutically acceptable acid. When a compound for use according to the present invention, contains a free acid such salts are prepared in a conventional manner by treating a solution or suspension of the compound with a chemical equivalent of a pharmaceutically acceptable base. Physiologically acceptable salts of a compound with a hydroxy group include the anion of said compound in combination with a suitable cation such as sodium or ammonium ion. Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds for use according to the present invention and these form a further aspect of the present invention.

[00278] For parenteral administration, solutions of the present compounds in sterile aqueous solution, aqueous propylene glycol or sesame or peanut oil may be employed. Such aqueous solutions should be suitable buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

[00279] Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. Examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, peanut oil, olive oil, syrup, phospholipids, gelatine, lactose, terra alba, sucrose, cyclodextrin, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, silicic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, polyoxyethylene, hydroxymethylcellulose and polyvinylpyrrolidone.

Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavouring agents.

[00280] The pharmaceutical compositions formed by combining the compounds of the invention and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

[00281] Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and which may include a suitable excipient. These formulations may be in the form of powder or granules, as a solution or suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion.

[00282] Compositions intended for oral use may be prepared according to any known method, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents, and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain the active ingredient in admixture with non-toxic pharmaceutically-acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch or alginic acid; binding agents, for example, starch, gelatine or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in U.S. Patent Nos. 4,356,108; 4,166,452; and 4,265,874, incorporated herein by reference, to form osmotic therapeutic tablets for controlled release.

[00283] Formulations for oral use may also be presented as hard gelatine capsules where the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or a soft gelatine capsule wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

[00284] Aqueous suspensions may contain the active compounds in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide such as lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethylenoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more

colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose or saccharin.

[00285] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as a liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

[00286] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active compound in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavouring, and colouring agents may also be present.

[00287] The pharmaceutical compositions comprising a compound for use according to the present invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example a liquid paraffin, or a mixture thereof. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

[00288] Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, preservative and flavouring and colouring agent. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known methods using suitable dispersing or wetting agents and suspending agents described above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conveniently employed as solvent or suspending medium. For this purpose, any bland fixed oil may be employed using synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[00289] The compositions may also be in the form of suppositories for rectal administration of the compounds of the present invention. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will thus melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols, for example.

[00290] For topical use, creams, ointments, jellies, solutions of suspensions, etc., containing the compounds of the present invention are contemplated. For the purpose of this application, topical applications shall include mouth washes and gargles.

[00291] The compounds for use according to the present invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes may be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

[00292] In addition, some of the compounds for use according to the present invention may form solvates with water or common organic solvents. Such solvates are also encompassed within the scope of the present invention.

[00293] Thus, in a further embodiment, there is provided a pharmaceutical composition comprising a compound for use according to the present invention, or a pharmaceutically acceptable salt, solvate, or prodrug thereof, and one or more pharmaceutically acceptable carriers, excipients, or diluents.

[00294] If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

[00295] A typical tablet which may be prepared by conventional tableting techniques may contain:

[00296] Core:

[00297] Active compound (as free compound or salt thereof) 5.0 mg

[00298] Lactosum PH. Eur. 67.8 mg

[00299] Cellulose, microcryst. (Avicel) 31.4 mg

[00300] Amberlite®IRP88* 1.0 mg

[00301] Magnesii stearas PH. Eur. q.s.

[00302] Coating:

[00303]	Hydroxypropyl methylcellulose	approx. 9 mg
[00304]	Mywacett 9-40 T**	approx. 0.9 mg

[00305] Polacrillin potassium NF, tablet disintegrant, Rohm and Haas.

5 [00306] ** Acylated monoglyceride used as plasticizer for film coating.

[00307] The compounds of the invention may be administered to a patient which is a mammal, especially a human in need thereof. Such mammals include also animals, both domestic animals, e.g., household pets, and non-domestic animals such as wildlife.

10 [00308] Any novel feature or combination of features described herein is considered essential to this invention.

[00309] The present invention also relate to the below methods of preparing the compounds of the invention.

[00310] The present invention is further illustrated in the following representative examples
15 which are, however, not intended to limit the scope of the invention in any way.

EXAMPLES

[00311] The following examples and general procedures refer to intermediate compounds and final products for general formula (I) identified above. The preparation of the compounds
20 of general formula (I) of the present invention is described in detail using the following examples. Occasionally, the reaction may not be applicable as described to each compound included within the disclosed scope of the invention. The compounds for which this occurs will be readily recognised by those skilled in the art. In these cases the reactions can be successfully performed by conventional modifications known to those skilled in the art, which is,
25 by appropriate protection of interfering groups, by changing to other conventional reagents, or by routine modification of reaction conditions. Alternatively, other reactions disclosed herein or otherwise conventional will be applicable to the preparation of the corresponding compounds of the invention. In all preparative methods, all starting materials are known or may easily be prepared from known starting materials. The structures of the compounds are
30 confirmed by either elemental analysis or nuclear magnetic resonance (NMR), where peaks assigned to characteristic protons in the title compounds are presented where appropriate. ¹H NMR shifts (δ_H) are given in parts per million (ppm) down field from tetramethylsilane as internal reference standard. M.p.: is melting point and is given in °C and is not corrected. Column chromatography was carried out using the technique described by W.C. Still *et al.*, *J. Org. Chem.* 43: 2923 (1978) on Merck silica gel 60 (Art. 9385). HPLC analyses are per-
35

formed using 5 μ m C18 4 x 250 mm column eluted with various mixtures of water and acetonitrile, flow = 1 ml/min, as described in the experimental section.

[00312] Microwave oven synthesis: The reaction was heated by microwave irradiation in sealed microwave vessels in a single mode Emrys Optimizer EXP from PersonalChemistry®.

- 5 **[00313]** Preparative HPLC: Column: 1.9 x 15 cm Waters XTerra RP-18. Buffer: linear gradient 5 – 95 % in 15 min, MeCN, 0.1 % TFA, flow rate of 15 ml/min. The pooled fractions are either evaporated to dryness *in vacuo*, or evaporated *in vacuo* until the MeCN is removed, and then frozen and freeze dried.

10 **[00314]** ABBREVIATIONS

[00315] d=day(s)

[00316] g=gram(s)

[00317] h=hour(s)

[00318] Hz=hertz

- 15 **[00319]** L=liter(s)

[00320] M=molar

[00321] mbar=millibar

[00322] mg=milligram(s)

[00323] min=minute(s)

- 20 **[00324]** mL=milliliter(s)

[00325] mM=millimolar

[00326] mmol=millimole(s)

[00327] mol=mole(s)

[00328] N=normal

- 25 **[00329]** ppm=parts per million

[00330] ESI=electrospray ionization

[00331] i.v.=intravenous

[00332] m/z=mass to charge ratio

[00333] MS=mass spectrometry

- 30 **[00334]** HPLC=high pressure liquid chromatography

[00335] RP=reverse phase

[00336] HPLC-MS=high pressure liquid chromatography - mass spectrometry

[00337] NMR=nuclear magnetic resonance spectroscopy

[00338] p.o.=per oral

- 35 **[00339]** rt=room temperature

[00340] s.c.=subcutaneous

[00341] t_r =retention time

[00342] DCM=dichloromethane, CH_2Cl_2 , methylenechloride

[00343] DIPEA=*N,N*-diisopropylethylamine

5 [00344] DMF=*N,N*-dimethylformamide

[00345] DMSO=dimethylsulfoxide

[00346] EDAC=1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride

[00347] Et_2O =diethyl ether

[00348] EtOAc=ethyl acetate

10 [00349] HOBt=1-hydroxybenzotriazole

[00350] MeCN=acetonitrile

[00351] MeOH=methanol

[00352] NMP=*N*-methylpyrrolidin-2-one

[00353] THF=tetrahydrofuran

15 [00354] CDCl_3 =deuterio chloroform

[00355] CD_3OD =tetra deuterio methanol

[00356] $\text{DMSO}-d_6$ = hexadeuterio dimethylsulfoxide

[00357] ANALYSIS

20 [00358] NMR

[00359] NMR spectra were recorded at 300 and 400 MHz on a Bruker DRX300, DRX400 or AV400 instrument equipped with 5 mm selective-inverse (SEI, ^1H and ^{13}C), 5 mm broad-band inverse (BBI, ^1H , broad-band) and 5 mm quadro nuclear (QNP, ^1H , ^{13}C) probeheads, respectively. Shifts (δ) are given in parts per million (ppm) down field from tetramethylsilane as internal reference standard.

25

[00360] PREPARATIVE TECHNIQUES

[00361] HPLC method

[00362] The RP-purification was performed on a Gilson system (3 Gilson 306 pumps, Gilson 170 DAD detector and a Gilson 215 liquidhandler) using a Waters X-terra RP (10 μm , 10 mm x 150 mm) with gradient elution, 5 % to 95 % solvent B (0.05 % TFA in acetonitrile) in solvent A (0.05 % TFA in water) within 15 min, 15 mL/min, detection at 210 nm, temperature rt. The pooled fractions are either evaporated to dryness *in vacuo*, or evaporated *in vacuo* until the acetonitrile is removed, and then frozen and freeze dried.

30


[00363] GENERAL

[00364] The following examples and general procedures refer to intermediate compounds and final products for general formula (I) identified in the specification and in the synthesis schemes. The preparation of the compounds of general formula (I) of the present invention is described in detail using the following examples. Occasionally, the reaction may not be applicable as described to each compound included within the disclosed scope of the invention. The compounds for which this occurs will be readily recognised by those skilled in the art. In these cases the reactions can be successfully performed by conventional modifications known to those skilled in the art, which is, by appropriate protection of interfering groups, by changing to other conventional reagents, or by routine modification of reaction conditions. Alternatively, other reactions disclosed herein or otherwise conventional will be applicable to the preparation of the corresponding compounds of the invention. In all preparative methods, all starting materials are known or may be prepared by a person skilled in the art in analogy with the preparation of similar known compounds or by the General procedure A described herein. The structures of the compounds are confirmed by either elemental analysis or nuclear magnetic resonance (NMR), where peaks assigned to characteristic protons in the title compounds are presented where appropriate.

[00365] GENERAL PROCEDURES

[00366] General procedure (A)


[00367] Compounds of the formula (Ia) according to the invention wherein R^1 , R^3 and R^4 are as defined for formula (I), with NR^3R^4 corresponding to ring A, can be prepared as outlined below:



[00368] An indole of formula (II) wherein NR^3R^4 is ring A as defined above may be reacted with an alkyl/benzyl halide of the formula (III) wherein R^1 is as defined above. This reaction may be carried out in a suitable solvent (e.g., DMF) at a temperature of up to reflux.

[00369] NNC 0####-0000-####-Example 1


3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester



[00370] To a solution of (1*H*-indol-4-yl)-(1,3,3-trimethyl-6-aza-bicyclo[3.2.1]oct-6-yl)-methanone (82 mg, 0.27 mmol) in dry DMF (2 mL) at room temperature under an inert atmosphere of nitrogen was added sodium hydride (17 mg, 0.69 mmol, 60% dispersion in oil), and after stirring for 30 min ethyl bromopropionate (100 mg, 0.55 mmol) was added, and the reaction mixture was stirred for 72 h at 60°C. The reaction was quenched by the addition of water (20 mL) followed by extraction with DCM (3 x 50 mL). The combined organic phases were dried (MgSO₄), filtered, and evaporated *in vacuo*. The resulting solid was purified by preparative HPLC, dried *in vacuo* at 50°C affording 29 mg (27%) of the title compound as a solid. MS-ESI *m/z* 397; ¹H NMR (400 MHz, DMSO) δ 0.88-1.08 (m, 6H), 1.14-1.22 (m, 7H), 1.28-1.29 (m) and 1.58-1.67 (m, 1H), 1.37-1.43 (m, 3H), 1.77-1.81 (m, 1H), 2.32-2.36 (m) and 3.10-3.20 (m, 1H), 2.81 (t, 2H), 3.35 (dd) and 3.75 (dd, 1H), 4.02-4.04 (m) and 4.68-4.70 (m, 1H), 4.11 (q, 2H), 4.46 (t, 2H), 6.44-6.50 (m, 1H), 7.10-7.13 (m, 1H), 7.17-7.26 (m, 2H), 7.38 (d, 1H).

[00371] NNC 0###-0000-####-Example 2

[00372] 3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid



[00373] To a solution of 3-[4-(1,3,3-trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester (25 mg, 0.063 mmol) in ethanol (1 mL) was added 1N aqueous sodium hydroxide solution (1 mL), and the resulting solution stirred for 2 h. The solution was acidified with 1N hydrochloric acid solution before the organic volatiles were removed *in vacuo* and the aqueous residues were extracted with DCM (3 x 5 mL). The combined organic phases were dried (MgSO₄), filtered, and evaporated *in vacuo*. The resulting solid was purified by preparative HPLC, dried *in vacuo* at 50°C affording 10 mg (43%) of the title compound as a solid. MS-ESI *m/z* 369; ¹H NMR (400 MHz, DMSO) δ 0.81-0.99 (m, 6H), 1.08-1.10 (m, 4H), 1.18-1.22 (m) and 1.50-1.54 (m, 1H), 1.30-1.36 (m, 3H), 1.71-1.73 (m, 1H), 2.24-2.28 (m) and 3.02-3.12 (m, 1H), 2.65 (t, 2H), 3.29 (dd) and 3.68 (dd, 1H), 3.93-3.95 (m)

and 4.60-4.62 (m, 1H), 4.29 (t, 2H), 6.24-6.29 (m, 1H), 6.94-6.96 (m, 1H), 7.00-7.12 (m, 2H), 7.27 (d, 1H).

[00374] PHARMACOLOGICAL METHODS

5 **[00375]** 11 β HSD1 enzyme assay

[00376] Materials

[00377] 3 H-cortisone and anti-rabbit Ig coated scintillation proximity assay (SPA) beads were purchased from Amersham Pharmacia Biotech, β -NADPH was from Sigma and rabbit anti-cortisol antibodies were from Fitzgerald. An extract of yeast transformed with h-11 β HSD1
10 (Hult et al., FEBS Lett., 441, 25 (1998)) was used as the source of enzyme. The test compounds were dissolved in DMSO (10 mM). All dilutions were performed in a buffer containing 50 mM TRIS-HCl (Sigma Chemical Co), 4 mM EDTA (Sigma Chemical Co), 0.1% BSA (Sigma Chemical Co), 0.01% Tween-20 (Sigma Chemical Co) and 0.005% bacitracin (Novo Nordisk A/S), pH=7.4. Optiplate 96 wells plates were supplied by Packard. The amount of
15 3 H-cortisol bound to the SPA beads was measured on TopCount NXT, Packard.

[00378] Methods


[00379] h-11 β HSD1, 120 nM 3 H-cortisone, 4 mM β -NADPH, antibody (1:200), serial dilutions of test compound and SPA particles (2 mg/well) were added to the wells. The reaction was initiated by mixing the different components and was allowed to proceed under shaking for
20 60 min at 30°C. The reaction was stopped by the addition of 10 fold excess of a stopping buffer containing 500 μ M carbenoxolone and 1 μ M cortisone. Data was analysed using GraphPad Prism software.

[00380] While the invention has been described and illustrated with reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications, and substitutions can be made therein without departing from the spirit and
25 scope of the present invention. For example, effective dosages other than the preferred dosages as set forth herein may be applicable as a consequence of variations in the responsiveness of the mammal being treated for the disease(s). Likewise, the specific pharmacological responses observed may vary according to and depending on the particular active
30 compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. Accordingly, the invention is not to be limited as by the appended claims.

[00381] The features disclosed in the foregoing description and/or in the claims may both separately and in any combination thereof be material for realising the invention in diverse forms thereof.

[00382] Preferred features of the invention:

- 5 1. Use of a substituted amide, a prodrug thereof, or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture or any tautomeric forms, wherein the substituted amide or a prodrug thereof is of formula I:




wherein:

X is selected from CR⁵ and N;

- 15 R¹ is selected from H and C₁-C₆alkyl-R⁶, wherein the alkyl group is substituted with 0-3 R⁷;

R² is selected from hydrogen, halo, C₁-C₆alkyl, and -C(=O)R¹³;

alternatively, R¹ and R² are, independently,



Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulphur;

- 25 Ring A is substituted with 0-3 groups selected from C₁-C₈alkyl, halo, hydroxy, oxo, cyano, C₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkylene, and C₁-C₆alkylcarbonyl, wherein each alkyl/alkylene group is substituted with 0-3 R¹⁴;

R⁵ is selected from hydrogen, C₁-C₆alkyl, -C(=O)R¹³, and cyano;

R⁶ is selected from cyano, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹,
 5 -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₁₀cycloalkyl, -N(R¹⁸)C(=O)-3-10 membered hetcycloalkyl, -N(R¹⁸)C(=O)-aryl, -N(R¹⁸)C(=O)-hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R¹⁶;

R⁷ is selected from halo, hydroxy, oxo, cyano, and C₁-C₈alkyl;
 10

R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from hydrogen, C₁-C₈alkyl, F, trihalomethyl, trihalomethoxy, hydroxy, and C₁-C₆alkyloxy, wherein the C₁-C₈alkyl and C₁-C₆alkyloxy are substituted with 0-3 R¹⁷;

15 alternatively, R⁸ and R⁹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy,
 20 arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

alternatively, R¹⁰ and R¹¹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein
 25 this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

alternatively, R⁸ and R¹⁰ together with the two carbon atoms to which they are attached form
 30 a saturated or partially saturated ring consisting of the two shown carbon atoms, 1-4 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy, and hetarylC₁-C₆alkyloxy;

35

R^{12} is selected from H, OH, $NR^{18}R^{19}$, C_3 - C_{10} cycloalkyl, 3-10 membered heterocycloalkyl, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, and $-C(=NR^{15})NR^{16}$; wherein the cycloalkyl and heterocycloalkyl groups are substituted with 0-3 R^{17} ;

- 5 R^{13} is selected from OH, C_1 - C_8 alkyl, C_1 - C_8 alkyloxy, C_1 - C_8 alkyloxy C_1 - C_4 alkylene, aryl, hetaryl, aryloxy, heteraryloxy, and $NR^{18}R^{19}$;

R^{14} is selected from halo, hydroxy, oxo, and cyano;

- 10 R^{15} and R^{16} are independently selected from H, C_1 - C_8 alkyl, 3-10 membered cycloalkyl, halo, OH, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-3 R^{20} ;

- 15 R^{17} is selected from halo, OH, oxo, nitro, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, $NR^{18}R^{19}$, C_1 - C_8 alkyl, C_1 - C_8 alkyloxy, and aryloxy;

R^{18} and R^{19} are independently selected from H, C_1 - C_8 alkyl, C_1 - C_8 alkyloxy, aryl, hetaryl, aryl C_1 - C_6 alkylene, and hetaryl C_1 - C_6 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-3 R^{20} ;

20

alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-3 C_1 - C_8 alkyl, aryl, hetaryl, aryl C_1 - C_6 alkylene, hetaryl C_1 - C_6 alkylene, hydroxy, oxo, C_1 - C_6 alkyloxy, aryl C_1 - C_6 alkyloxy, hetaryl C_1 - C_6 alkyloxy, C_1 - C_6 alkyloxy C_1 - C_6 alkyl, C_1 - C_6 alkylcarbonyl, arylcarbonyl, hetarylcarbonyl, aryl C_1 - C_6 alkylcarbonyl, hetaryl C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarboxy, arylcarboxy, hetarylcarboxy, aryl C_1 - C_6 alkylcarboxy, and hetaryl C_1 - C_6 alkylcarboxy;

25

- 30 R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_6 alkyl, C_1 - C_6 alkyloxy, $NR^{21}R^{22}$, methylenedioxy, dihalomethylenedioxy, trihalomethyl, and trihalomethyloxy;

R^{21} and R^{22} are independently selected from H, C_1 - C_8 alkyl, and aryl C_1 - C_6 alkyl;

35

R^{23} is selected from H and C_1 - C_6 alkyl;

n is selected from 0, 1, and 2;

Y is selected from O and S;


5

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

2. The use according to clause 1, wherein:

10 R^1 is selected from H and C_1 - C_4 alkyl- R^6 , wherein the alkyl group is substituted with 0-1 R^7 ;

R^2 is selected from hydrogen, C_1 - C_6 alkyl, and $-C(=O)R^{13}$;



alternatively, R^1 and R^2 are, independently,

15

Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen and 7-10 carbon atoms;

20 Ring A is substituted with 0-3 groups selected from C_1 - C_4 alkyl, halo, hydroxy, and C_1 - C_6 alkyloxy;

R^5 is selected from hydrogen and C_1 - C_4 alkyl;

25 R^6 is selected from cyano, C_3 - C_6 cycloalkyl, 3-6 membered hetcycloalkyl, aryl, hetaryl, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $-S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, $-N(R^{23})C(=Y)NR^{18}R^{19}$, $-C(=NR^{15})NR^{15}$, $-N(R^{18})C(=O)R^{13}$, $-N(R^{18})C(=O)-C_3$ - C_6 cycloalkyl, $-N(R^{18})C(=O)$ -3-6 membered hetcycloalkyl, $-N(R^{18})C(=O)$ -aryl, $-N(R^{18})C(=O)$ -hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R^{16} ;

30 R^7 is selected from halo and C_1 - C_4 alkyl;

R^8 , R^9 , R^{10} and R^{11} are each independently selected from hydrogen and C_1 - C_4 alkyl;

alternatively, R^8 and R^{10} together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms and 1-4 additional carbon atoms, wherein this ring is substituted with 0-1 groups selected from halo, trihalomethyl, hydroxyl, and C_1 - C_6 alkyl;

5

R^{12} is selected from H, OH, and $NR^{18}R^{19}$;

R^{13} is selected from OH, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, C_1 - C_4 alkyloxy C_1 - C_4 alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and $NR^{18}R^{19}$;

10

R^{15} and R^{16} are independently selected from H, C_1 - C_4 alkyl, 3-6 membered cycloalkyl, halo, OH, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-1 R^{20} ;

15

R^{18} and R^{19} are independently selected from H, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, aryl, hetaryl, aryl C_1 - C_4 alkylene, and hetaryl C_1 - C_4 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-1 R^{20} ;

20

alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-5 carbon atoms, and 0-1 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-1 C_1 - C_4 alkyl, aryl, hetaryl, aryl C_1 - C_4 alkylene, hetaryl C_1 - C_4 alkylene, hydroxy, and C_1 - C_4 alkyloxy;

25

R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, $NR^{21}R^{22}$, trihalomethyl, and trihalomethyloxy;

R^{21} and R^{22} are independently selected from H, C_1 - C_4 alkyl, and aryl C_1 - C_4 alkyl;

30


R^{23} is selected from H and C_1 - C_6 alkyl;

n is selected from 0, 1, and 2; and,

Y is selected from O and S.


35

3. The use according to clause 1, wherein the substituted amide or prodrug thereof is of formula IA:




IA.

4. The use according to clause 1, wherein the substituted amide or prodrug thereof is of formula IB:




IB.

5. The use according to clause 1, wherein the substituted amide or prodrug thereof is of formula IC:



IC.


6. The use according to clause 1, wherein the substituted amide or prodrug thereof is of formula ID:



5

7. The use according to clause 1, wherein:


Ring A is selected from:



- 10 Ring A is substituted with 0-2 R^{24} ; and,

R^{24} is selected from C_1 - C_8 alkyl, halo, hydroxy, oxo, cyano, and C_1 - C_6 alkyloxy.

- 15 8. The use according to clause 7, wherein:



- 20 9. The use according to clause 1, wherein the substituted amide or a prodrug thereof is of the selected from the group:

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester;

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid;
or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or
5 mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

10. The use according to any of clauses 1-9, for the preparation of a pharmaceutical composition for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

11. The use according to clause 10, wherein the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.

12. The use according to clause 10, wherein the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), the progression from IGT to type 2 diabetes, the progression of the metabolic syndrome into type 2 diabetes, diabetic late complications, neurodegenerative and psychiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

13. The use according to any of the clauses 10-12, wherein the pharmaceutical composition is suitable for a route of administration selected from oral, nasal, buccal, transdermal, pulmonary, and parenteral.

14. A method for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial, the method comprising administering to a subject in need thereof an effective amount of a compound according to any of clauses 1-9.

15. The method according to clause 14, wherein the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.


16. The method according to clause 14, wherein the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity,

type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), progression from IGT to type 2 diabetes, progression of metabolic syndrome into type 2 diabetes, diabetic late complications, neurodegenerative and psychiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

5

17. The method according to any of clauses 14-16, wherein the administering is via a route selected from oral, nasal, buccal, transdermal, pulmonal, and parenteral.

18. A compound of the formula I:



10

wherein:

X is selected from CR⁵ and N;


15

R¹ is selected from H and C₁-C₆alkyl-R⁶, wherein the alkyl group is substituted with 0-3 R⁷;

R² is selected from hydrogen, halo, C₁-C₆alkyl, and -C(=O)R¹³;

20

alternatively, R¹ and R² are, independently,



Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulphur;

25

Ring A is substituted with 0-3 groups selected from C₁-C₈alkyl, halo, hydroxy, oxo, cyano, C₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkylene, and C₁-C₆alkylcarbonyl, wherein each alkyl/alkylene group is substituted with 0-3 R¹⁴;

R⁵ is selected from hydrogen, C₁-C₆alkyl, -C(=O)R¹³, and cyano;

5 R⁶ is selected from cyano, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹, -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₁₀cycloalkyl, -N(R¹⁸)C(=O)-3-10 membered hetcycloalkyl, -N(R¹⁸)C(=O)-aryl, -N(R¹⁸)C(=O)-hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R¹⁶;

10 R⁷ is selected from halo, hydroxy, oxo, cyano, and C₁-C₈alkyl;

R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from hydrogen, C₁-C₈alkyl, F, trihalomethyl, trihalomethoxy, hydroxy, and C₁-C₆alkyloxy, wherein the C₁-C₈alkyl and C₁-C₆alkyloxy are substituted with 0-3 R¹⁷;

15

alternatively, R⁸ and R⁹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, 20 hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

alternatively, R¹⁰ and R¹¹ together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional 25 carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, arylC₁-C₆alkyloxy or hetarylC₁-C₆alkyloxy;

30 alternatively, R⁸ and R¹⁰ together with the two carbon atoms to which they are attached form a saturated or partially saturated ring consisting of the two shown carbon atoms, 1-4 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C₁-C₆alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, aryloxy, 35 arylC₁-C₆alkyloxy, and hetarylC₁-C₆alkyloxy;

R¹² is selected from H, OH, NR¹⁸R¹⁹, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, and -C(=NR¹⁵)NR¹⁶; wherein the cycloalkyl and hetcycloalkyl groups are substituted with 0-3 R¹⁷;

5

R¹³ is selected from OH, C₁-C₈alkyl, C₁-C₈alkyloxy, C₁-C₈alkyloxyC₁-C₄alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and NR¹⁸R¹⁹;

R¹⁴ is selected from halo, hydroxy, oxo, and cyano;

10

R¹⁵ and R¹⁶ are independently selected from H, C₁-C₈alkyl, 3-10 membered cycloalkyl, halo, OH, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-3 R²⁰;

15

R¹⁷ is selected from halo, OH, oxo, nitro, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, NR¹⁸R¹⁹, C₁-C₈alkyl, C₁-C₆alkyloxy, and aryloxy;

R¹⁸ and R¹⁹ are independently selected from H, C₁-C₈alkyl, C₁-C₈alkyloxy, aryl, hetaryl, arylC₁-C₆alkylene, and hetarylC₁-C₆alkylene, wherein the alkyl/alkylene, aryl, and hetaryl groups are independently substituted with 0-3 R²⁰;

20

alternatively, R¹⁸ and R¹⁹, together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-3 C₁-C₈alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, arylC₁-C₆alkyloxy, hetarylC₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkyl, C₁-C₆alkylcarbonyl, arylcarbonyl, hetarylcarbonyl, arylC₁-C₆alkylcarbonyl, hetarylC₁-C₆alkylcarbonyl, C₁-C₆alkylcarboxy, arylcarboxy, hetarylcarboxy, arylC₁-C₆alkylcarboxy, and hetarylC₁-C₆alkylcarboxy;

30

R²⁰ is selected from H, OH, oxo, halo, cyano, nitro, C₁-C₆alkyl, C₁-C₆alkyloxy, NR²¹R²², methylenedioxy, dihalomethylenedioxy, trihalomethyl, and trihalomethyloxy;

R²¹ and R²² are independently selected from H, C₁-C₈alkyl, and arylC₁-C₆alkyl;

35

R^{23} is selected from H and C_1 - C_6 alkyl;

n is selected from 0, 1, and 2;

5 Y is selected from O and S;


or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

10

19. The compound according to clause 18, wherein:

R^1 is selected from H and C_1 - C_4 alkyl- R^6 , wherein the alkyl group is substituted with 0-1 R^7 ;

15 R^2 is selected from hydrogen, C_1 - C_6 alkyl, and $-C(=O)R^{13}$;



alternatively, R^1 and R^2 are, independently,

20

Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen and 7-10 carbon atoms;

Ring A is substituted with 0-3 groups selected from C_1 - C_4 alkyl, halo, hydroxy, and C_1 - C_6 alkyloxy;

25

R^5 is selected from hydrogen and C_1 - C_4 alkyl;

30

R^6 is selected from cyano, C_3 - C_6 cycloalkyl, 3-6 membered hetcycloalkyl, aryl, hetaryl, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $-S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, $-N(R^{23})C(=Y)NR^{18}R^{19}$, $-C(=NR^{15})NR^{15}$, $-N(R^{18})C(=O)R^{13}$, $-N(R^{18})C(=O)-C_3$ - C_6 cycloalkyl, $-N(R^{18})C(=O)-3$ -6 membered hetcycloalkyl, $-N(R^{18})C(=O)$ -aryl, $-N(R^{18})C(=O)$ -hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R^{16} ;

R^7 is selected from halo and C_1 - C_4 alkyl;

R^8 , R^9 , R^{10} and R^{11} are each independently selected from hydrogen and C_1 - C_4 alkyl;

alternatively, R^8 and R^{10} together with the two carbon atoms to which they are attached form
5 a saturated or partially saturated ring consisting of the two shown carbon atoms and 1-4
additional carbon atoms, wherein this ring is substituted with 0-1 groups selected from halo,
trihalomethyl, hydroxyl, and C_1 - C_6 alkyl;

R^{12} is selected from H, OH, and $NR^{18}R^{19}$;

R^{13} is selected from OH, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, C_1 - C_4 alkyloxy C_1 - C_4 alkylene, aryl,
10 hetaryl, aryloxy, hetaryloxy, and $NR^{18}R^{19}$;

R^{15} and R^{16} are independently selected from H, C_1 - C_4 alkyl, 3-6 membered cycloalkyl, halo,
15 OH, cyano, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, aryl, and hetaryl,
wherein the alkyl and cycloalkyl groups are substituted with 0-1 R^{20} ;

R^{18} and R^{19} are independently selected from H, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, aryl, hetaryl,
20 aryl C_1 - C_4 alkylene, and hetaryl C_1 - C_4 alkylene, wherein the alkyl/alkylene, aryl, and hetaryl
groups are independently substituted with 0-1 R^{20} ;

alternatively, R^{18} and R^{19} , together with the nitrogen atom to which they are attached, form
a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the
shown nitrogen, 4-5 carbon atoms, and 0-1 additional heteroatoms selected from nitrogen,
25 oxygen, and sulfur, wherein this ring is substituted with 0-1 C_1 - C_4 alkyl, aryl, hetaryl, aryl C_1 -
 C_4 alkylene, hetaryl C_1 - C_4 alkylene, hydroxy, and C_1 - C_4 alkyloxy;

R^{20} is selected from H, OH, oxo, halo, cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 alkyloxy, $NR^{21}R^{22}$, tri-
30 halomethyl, and trihalomethoxy;


R^{21} and R^{22} are independently selected from H, C_1 - C_4 alkyl, and aryl C_1 - C_4 alkyl;

R^{23} is selected from H and C_1 - C_6 alkyl;


n is selected from 0, 1, and 2; and,

Y is selected from O and S.


- 5 20. The compound according to clause 18, wherein the compound is of formula IA:



- 10 21. The compound according to clause 18, wherein the compound is of formula IB:




- 15 22. The compound according to clause 18, wherein the compound is of formula IC:



62


23. The compound according to clause 18, wherein the compound is of formula ID:



5

24. The compound according to clause 18, wherein:

Ring A is selected from:




10

Ring A is substituted with 0-2 R^{24} ; and,

R^{24} is selected from C_1 - C_8 alkyl, halo, hydroxy, oxo, cyano, and C_1 - C_6 alkyloxy.

15

25. The use according to clause 24, wherein:



20

26. The compound according to clause 18, wherein the compound is selected from the group:

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester;

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid; or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

27. A compound according to any one of the clauses 18-26, which is an agent useful for the treatment of conditions, disorders, or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

28. The compound according to clause 27, wherein the conditions, disorders, and diseases that are influenced by intracellular glucocorticoid levels.


29. The compound according to clause 27, wherein the conditions, disorders, or diseases are selected from metabolic syndrome, insulin resistance, dyslipidemia, hypertension, obesity, type 2 diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), progression from IGT to type 2 diabetes, progression of metabolic syndrome into type 2 diabetes, diabetic late complications, neurodegenerative and psychiatric disorders, and the adverse effects of glucocorticoid receptor agonist treatment or therapy.

30. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of the clauses 18-27 together with one or more pharmaceutically acceptable carriers or excipients.

31. The pharmaceutical composition according to clause 30 which is suitable for oral, nasal, buccal, transdermal, pulmonary, or parenteral administration.

WHAT IS CLAIMED:

1. A compound of the formula I:




wherein:

X is selected from CR⁵ and N;

R¹ is selected from H and C₁-C₆alkyl-R⁶, wherein the alkyl group is substituted with 0-3 R⁷;

R² is selected from hydrogen, halo, C₁-C₆alkyl, and -C(=O)R¹³;



alternatively, R¹ and R² are, independently,

Ring A is a saturated or partially saturated bicyclic or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulphur;

Ring A is substituted with 0-3 groups selected from C₁-C₈alkyl, halo, hydroxy, oxo, cyano, C₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkylene, and C₁-C₆alkylcarbonyl, wherein each alkyl/alkylene group is substituted with 0-3 R¹⁴;

R⁵ is selected from hydrogen, C₁-C₆alkyl, -C(=O)R¹³, and cyano;

R⁶ is selected from cyano, C₃-C₁₀cycloalkyl, 3-10 membered hetcycloalkyl, aryl, hetaryl, -C(=O)R¹³, -S(=O)_nR¹³, -S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, -N(R²³)C(=Y)NR¹⁸R¹⁹, -C(=NR¹⁵)NR¹⁵, -N(R¹⁸)C(=O)R¹³, -N(R¹⁸)C(=O)-C₃-C₁₀cycloalkyl, -N(R¹⁸)C(=O)-3-10 mem-

bered hetcycloalkyl, $-N(R^{18})C(=O)$ -aryl, $-N(R^{18})C(=O)$ -hetaryl, wherein the cycloalkyl, hetcycloalkyl, aryl, and hetaryl groups are substituted with 0-3 R^{16} ;

R^7 is selected from halo, hydroxy, oxo, cyano, and C_1 - C_8 alkyl;

5

R^8 , R^9 , R^{10} and R^{11} are each independently selected from hydrogen, C_1 - C_8 alkyl, F, trihalomethyl, trihalomethoxy, hydroxy, and C_1 - C_6 alkyloxy, wherein the C_1 - C_8 alkyl and C_1 - C_6 alkyloxy are substituted with 0-3 R^{17} ;

- 10 alternatively, R^8 and R^9 together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C_1 - C_6 alkyl, aryl, hetaryl, aryl C_1 - C_6 alkylene, hetaryl C_1 - C_6 alkylene, hydroxy, oxo, C_1 - C_6 alkyloxy, aryloxy, aryl C_1 -
15 C_6 alkyloxy or hetaryl C_1 - C_6 alkyloxy;

- alternatively, R^{10} and R^{11} together with the carbon atom to which they are attached form a saturated or partially saturated ring consisting of the carbon atom shown, 2-5 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this
20 ring is substituted with 0-3 groups selected from halo, trihalomethyl, C_1 - C_6 alkyl, aryl, hetaryl, aryl C_1 - C_6 alkylene, hetaryl C_1 - C_6 alkylene, hydroxy, oxo, C_1 - C_6 alkyloxy, aryloxy, aryl C_1 - C_6 alkyloxy or hetaryl C_1 - C_6 alkyloxy;

- alternatively, R^8 and R^{10} together with the two carbon atoms to which they are attached form
25 a saturated or partially saturated ring consisting of the two shown carbon atoms, 1-4 additional carbon atoms, and 0-2 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein this ring is substituted with 0-3 groups selected from halo, trihalomethyl, C_1 - C_6 alkyl, aryl, hetaryl, aryl C_1 - C_6 alkylene, hetaryl C_1 - C_6 alkylene, hydroxy, oxo, C_1 - C_6 alkyloxy, aryloxy, aryl C_1 - C_6 alkyloxy, and hetaryl C_1 - C_6 alkyloxy;

30

R^{12} is selected from H, OH, $NR^{18}R^{19}$, C_3 - C_{10} cycloalkyl, 3-10 membered hetcycloalkyl, $-C(=O)R^{13}$, $-S(=O)_nR^{13}$, $S(=O)_nNR^{18}R^{19}$, $-N(R^{18})S(=O)_nR^{13}$, and $-C(=NR^{15})NR^{16}$; wherein the cycloalkyl and hetcycloalkyl groups are substituted with 0-3 R^{17} ;

R¹³ is selected from OH, C₁-C₈alkyl, C₁-C₈alkyloxy, C₁-C₈alkyloxyC₁-C₄alkylene, aryl, hetaryl, aryloxy, hetaryloxy, and NR¹⁸R¹⁹;

R¹⁴ is selected from halo, hydroxy, oxo, and cyano;

5

R¹⁵ and R¹⁶ are independently selected from H, C₁-C₈alkyl, 3-10 membered cycloalkyl, halo, OH, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, aryl, and hetaryl, wherein the alkyl and cycloalkyl groups are substituted with 0-3 R²⁰;

10 R¹⁷ is selected from halo, OH, oxo, nitro, cyano, -C(=O)R¹³, -S(=O)_nR¹³, S(=O)_nNR¹⁸R¹⁹, -N(R¹⁸)S(=O)_nR¹³, NR¹⁸R¹⁹, C₁-C₈alkyl, C₁-C₈alkyloxy, and aryloxy;

R¹⁸ and R¹⁹ are independently selected from H, C₁-C₈alkyl, C₁-C₈alkyloxy, aryl, hetaryl, arylC₁-C₆alkylene, and hetarylC₁-C₆alkylene, wherein the alkyl/alkylene, aryl, and hetaryl

15 groups are independently substituted with 0-3 R²⁰;

alternatively, R¹⁸ and R¹⁹, together with the nitrogen atom to which they are attached, form a saturated or partially saturated monocyclic, bicyclic, or tricyclic ring consisting of the shown nitrogen, 4-10 carbon atoms, and 0-2 additional heteroatoms selected from nitrogen, oxygen, and sulfur, wherein this ring is substituted with 0-3 C₁-C₈alkyl, aryl, hetaryl, arylC₁-C₆alkylene, hetarylC₁-C₆alkylene, hydroxy, oxo, C₁-C₆alkyloxy, arylC₁-C₆alkyloxy, hetarylC₁-C₆alkyloxy, C₁-C₆alkyloxyC₁-C₆alkyl, C₁-C₆alkylcarbonyl, arylcarbonyl, hetarylcarbonyl, arylC₁-C₆alkylcarbonyl, hetarylC₁-C₆alkylcarbonyl, C₁-C₆alkylcarboxy, arylcarboxy, hetarylcarboxy, arylC₁-C₆alkyl-carboxy, and hetarylC₁-C₆alkylcarboxy;

25

R²⁰ is selected from H, OH, oxo, halo, cyano, nitro, C₁-C₆alkyl, C₁-C₆alkyloxy, NR²¹R²², methylendioxo, dihalomethylendioxo, trihalomethyl, and trihalomethyloxy;

R²¹ and R²² are independently selected from H, C₁-C₈alkyl, and arylC₁-C₆alkyl;

30

R²³ is selected from H and C₁-C₆alkyl;


n is selected from 0, 1, and 2;

35 Y is selected from O and S;

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.


5

2. The compound according to claim 1, wherein the compound is of formula IA:




10

3. The compound according to claim 1 wherein the compound is of formula IB:




15

4. The compound according to claim 1, wherein the compound is of formula IC:



5. The compound according to claim 1, wherein the compound is of formula ID:



5

6. The compound according to claim 1, wherein the compound is selected from the group:
3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid ethyl ester;

3-[4-(1,3,3-Trimethyl-6-aza-bicyclo[3.2.1]octane-6-carbonyl)-indol-1-yl]-propionic acid, or
10 a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

7. The compound according to any one of the preceding claims, which is an agent useful for the treatment, prevention and/or prophylaxis of any conditions, disorders and diseases
15 wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

8. The compound according to any one of the claims 1-6, which is an agent useful for the treatment, prevention and/or prophylaxis of any conditions, disorders and diseases that are influenced by intracellular glucocorticoid levels.

20

9. The compound according to any one of the claims 1-6 which is an agent useful for the treatment, prevention and/or prophylaxis of conditions, disorders or diseases selected from the group consisting of the metabolic syndrome, insulin resistance, dyslipidemia, hypertension and obesity.

25

10. A pharmaceutical composition comprising, as an active ingredient, at least one compound according to any one of the claims 1-6 together with one or more pharmaceutically acceptable carriers or excipients.

11. Use of a compound according to any of the claims 1-6, for the preparation of a pharmaceutical composition for the treatment, prevention and/or prophylaxis of any conditions, disorders and diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial.

5

12. Use of a compound according to any of the claims 1-6, for the preparation of a pharmaceutical composition for the treatment, prevention and/or prophylaxis of any conditions, disorders and diseases that are influenced by intracellular glucocorticoid levels.

10

13. Use of a compound according to any of the claims 1-6, for the preparation of a pharmaceutical composition for the treatment, prevention and/or prophylaxis of conditions, disorders or diseases selected from the group consisting of the metabolic syndrome, insulin resistance, dyslipidemia, hypertension and obesity.

15

14. A method for the treatment, prevention and/or prophylaxis of any conditions, disorders or diseases wherein a modulation or an inhibition of the activity of 11 β HSD1 is beneficial, the method comprising administering to a subject in need thereof an effective amount of a compound according to the invention.

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2007/052219

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D403/06 A61K31/439 A61P3/10 A61P3/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2004/089470 A (NOVO NORDISK AS [DK]; ANDERSEN HENRIK SUNE [DK]; KAMPEN GITA CAMILLA T) 21 October 2004 (2004-10-21) claims page 68, line 20 - line 21 examples 4.6,4.33,6.3,6.4 -----	1-14
X	WO 2005/095397 A (ACTELION PHARMACEUTICALS LTD [CH]; FRETZ HEINZ [CH]; FECHER ANJA [CH];) 13 October 2005 (2005-10-13) claims example 149; table 8 -----	1,4,5, 7-10
X	WO 00/46197 A (ASTRAZENECA UK LTD [GB]; FAULL ALAN WELLINGTON [GB]; KETTLE JASON [GB]) 10 August 2000 (2000-08-10) claims example 8; table 1 -----	1,4,5, 7-10
-/--		

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

12 November 2007

Date of mailing of the international search report

23/11/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

KOLLMANNBERGER, M

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/052219

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/040329 A (NOVO NORDISK AS [DK]; KILBURN JOHN PAUL [DK]; ANDERSEN HENRIK SUNE [DK]) 20 April 2006 (2006-04-20) claims examples 1,2,2.2,2.3,2.7,2.10 -----	1-14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2007/052219

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 14 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers allsearchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search reportcovers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/052219

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004089470	A	21-10-2004	WO 2004089896 A1	21-10-2004
			EP 1618090 A1	25-01-2006
			EP 1615698 A2	18-01-2006
			JP 2006522746 T	05-10-2006
			JP 2006522748 T	05-10-2006
WO 2005095397	A	13-10-2005	AR 047928 A1	01-03-2006
			AU 2005229356 A1	13-10-2005
			CA 2558509 A1	13-10-2005
			CN 1930162 A	14-03-2007
			EP 1725553 A1	29-11-2006
			JP 2007526276 T	13-09-2007
			KR 20070029670 A	14-03-2007
WO 0046197	A	10-08-2000	AT 250577 T	15-10-2003
			AU 2304300 A	25-08-2000
			DE 60005485 D1	30-10-2003
			DE 60005485 T2	01-07-2004
			EP 1150953 A1	07-11-2001
			JP 2002536360 T	29-10-2002
			US 6613760 B1	02-09-2003
WO 2006040329	A	20-04-2006	EP 1802623 A1	04-07-2007